

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
13 June 2002 (13.06.2002)

PCT

(10) International Publication Number
WO 02/46369 A2

(51) International Patent Classification⁷: **C12N 1/19**,
15/81, 15/12, C12Q 1/02, 1/68, C07K 14/72, 14/705 //
G01N 33/68, C12R 1:645

SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN,
YU, ZA, ZM, ZW.

(21) International Application Number: PCT/GB01/05460

(22) International Filing Date:
10 December 2001 (10.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0030038.4 8 December 2000 (08.12.2000) GB

(71) Applicant (for all designated States except US): **SEPT-
GEN LIMITED** [GB/GB]; Barclays Venture Centre, Sir
William Lyons Road, Coventry CV4 7EZ (GB).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **DAVEY, John**
[GB/GB]; 132 Lyndon Road, Solihull, West Midlands B92
7QE (GB).

(74) Agents: **ELSY, David** et al.; Withers & Rogers, Goldings
House, 2 Hays Lane, London SE1 2HW (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Declaration under Rule 4.17:

— as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for the following des-
ignations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR,
BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ,
EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM,
PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR,
TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent
(GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB,
GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ,
CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG)

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: YEAST-BASED ASSAY

(57) Abstract: The application discloses *Sz. pombe* yeast cells which have been modified so that a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway is derepressed during the mitotic phase of cell growth. Isolated nucleic acid molecules encoding constructs used to make the yeast cells, uses of the cells and nucleic acid molecules to study GPCR pathways and to isolate compounds which effect in such pathways are also disclosed.

WO 02/46369 A2

Yeast-based Assay

The application relates to modified yeast cells which may be used to study the activity of G-protein coupled receptors (GPCRs). The yeast cells used are *Schizosaccharomyces pombe* (*Sz. pombe*) containing a reporter gene-promoter construct. The invention also relates to isolated nucleic acid encoding the reporter gene-promoter construct and to uses of the yeast cells and nucleic acid molecules in assays.

GPCRs are an important class of receptors in all eukaryotic organisms, including mammals and yeast, and are responsible for conveying hormonal and sensory signals to the cell machinery (reviewed in Baldwin, 1994). Such receptors have a common structure comprising 7-transmembrane domains with an extracellular N-terminus and C-terminal cytoplasmic tail. GPCRs are usually coupled to a heterotrimeric G protein composed of $G\alpha$, $G\beta$ and $G\gamma$ subunits. Binding of a ligand to the receptor stimulates a change in the G protein where guanosine diphosphate (GDP) bound to the $G\alpha$ subunit is exchanged for guanosine triphosphate (GTP). Accompanying conformational changes result in the dissociation of $G\alpha$ -GTP from the $G\beta\gamma$ dimer, either of which can modulate the activity of effector proteins to bring about changes in cell behaviour.

GPCRs control the physiology of all major organ systems and have been important targets for therapeutic and diagnostic advances, providing clinically successful drugs in nearly all the major pharmaceutical markets. Many of the 200 or so well characterised GPCRs are associated with at least one drug and about 60% of commercially available drugs act on GPCRs, providing some \$27 billion in annual sales world-wide. There are another 100 or so GPCRs for which ligands have not yet been identified. These so called 'orphan' receptors are likely to include many that will become important drug targets. Analysis of the human genome indicates that there are probably another 500 orphan GPCRs that will need to be characterised. There is therefore considerable interest in developing drug leads targeted at the GPCRs.

One approach to the identification of new drugs is the development of high throughput screens (HTS) for GPCRs. In most cases, the target GPCR is expressed in a host system

such that activation of the receptor leads to a change in cell behaviour. Screening can then identify drug leads that either mimic the action of the natural ligand (agonists) or block the receptor (antagonists). All eukaryotic cells contain GPCRs and each can be adapted for HTS but it is not always practical to do this and most screens use a limited range of host systems. These include mammalian cells, frog melanocytes, insect cells and yeast. Each system has its advantages and disadvantages. For example, mammalian cells might seem the obvious choice for studying human GPCRs but they are difficult and expensive to work with and screens are often complicated by the inherent presence of receptors closely related to the GPCR being studied. The presence of related receptors can also complicate screens involving frog melanocytes and insect cells. These problems do not apply to yeast and many have turned to using this relatively simple cell as a surrogate host for screening human GPCRs.

G-protein coupled receptors are known in yeast. Accordingly, yeast, such as *Saccharomyces cerevisiae* (*S. cerevisiae*) have been used to study GPCR-regulated signalling systems. Yeast cells are particularly advantageous because they have the ability to be easily manipulated, at low cost and with high levels of production. Unlike bacteria, yeast has the potential to perform eukaryotic post-translational modifications that may affect receptor function (Reiländer and Weib, 1998). The mechanism of transcriptional activation in yeast and higher eukaryotes may be very similar. For example, yeast upstream activation sites (UAS) and some transcriptional activators have been found to have very similar activity to that of their mammalian equivalents (Jones *et al.*, 1988).

Most work carried out on yeast systems has been on *S. cerevisiae*. There are many reports describing the coupling of exogenous GPCRs to the intracellular signalling machinery in *S. cerevisiae*. These include the human β_2 -adrenergic (King *et al.*, 1990), rat somatostatin (Price *et al.*, 1995; Bass *et al.*, 1996), rat adenosine A_{2A} (Price *et al.*, 1996), human growth hormone releasing hormone (Kajkowski *et al.*, 1997), human lysophosphatidic acid (Erickson *et al.*, 1998), human formyl peptide receptor like-1 (Klein *et al.*, 1998), human C5a chemoattractant (Klein *et al.*, 1998; Baranski *et al.*, 1999), mushroom pheromone (Olesnicki *et al.*, 1999), human somatostatin SST₂ (Brown *et al.*, 2000), human somatostatin SST₅ (Brown *et al.*, 2000), human serotonin 5-HT_{1A} (Brown *et al.*, 2000),

human serotonin 5-HT_{1D} (Brown *et al.*, 2000), human melatonin ML_{1B} (Brown *et al.*, 2000), human purinergic P2Y₁ (Brown *et al.*, 2000), human purinergic P2Y₂ (Brown *et al.*, 2000), human adenosine A_{2B} (Brown *et al.*, 2000), human UDP-glucose (Chambers *et al.*, 2000), human protease-activated receptor (Swift *et al.*, 2000), human muscarinic M₁ (Erlenbach *et al.*, 2001), human muscarinic M₃ (Erlenbach *et al.*, 2001), human muscarinic M₅ (Erlenbach *et al.*, 2001) and human vasopressin V₂ (Erlenbach *et al.*, 2001).

However, not all GPCRs couple to the signalling machinery in *S. cerevisiae*. Receptors that fail to couple are not normally reported but as many as 40% of human GPCRs are not functional in *S. cerevisiae*.

The fission yeast *Schizosaccharomyces pombe* (*Sz. pombe*) is becoming popular as an alternative genetically tractable eukaryote which is not only phylogenetically distant from *S. cerevisiae*, but in several aspects of its cell and molecular biology seems to more closely resemble a higher eukaryotic cell (Reiländer and Weib, 1998; Allshire *et al.*, 1987). *Sz. pombe* would therefore seem to provide an attractive alternative to the budding yeast. Unfortunately, all previously reported attempts to couple exogenous GPCRs to the signalling machinery in *Sz. pombe* have been unsuccessful. It appears that although the receptors are expressed they fail to couple to the intracellular signalling machinery in the yeast. Examples include bacteriorhodopsin (Hildebrandt *et al.*, 1993), human dopamine D_{2S} (Sander *et al.*, 1994), human neurokinin NK2 (Arkininstall *et al.*, 1995) and human β_2 -adrenergic (Ficca *et al.*, 1995).

This application describes how *Sz. pombe* may be manipulated, for example by way of modification, to allow the coupling of exogenous GPCRs to the intracellular signalling machinery and hence generate strains suitable for high throughput screening for agonists and antagonists that affect the activity of the exogenous receptors.

Fission yeasts, such as *Sz. pombe*, have two distinct growth cycles. Firstly, they have a normal vegetative or mitotic cycle in which haploid cells simply divide by fission. Secondly, they have a meiotic cycle. In the meiotic cycle, a yeast cell conjugates with a second yeast cell to form a diploid cell. The diploid cell then undergoes meiosis and

sporulation to form four haploid spores. Such spores are very resilient and the meiotic cycle is usually triggered when environmental conditions for the yeast no longer support mitotic growth. For example, the meiotic cycle in *Sz. pombe* is usually triggered by nitrogen starvation.

Conjugation in *Sz. pombe* is controlled by the reciprocal action of diffusible mating pheromones. M cells (of mating type minus) release M-factor which prepares P cells (mating type plus) for mating, while P cells release P-factor which stimulates M cells for mating. Binding of the pheromones to their receptors on the surface of the target cell activates an intracellular signalling pathway which leads to changes in the pattern of gene transcription and prepares the cell for mating. Responses induced by the pheromones include G₁ arrest of the cell cycle, an increase in agglutination, and the elongation of the cell to form a shmoo. The M-receptor and P-receptor to which the M-factor and P-factor pheromones bind are examples of G-protein coupled receptors. On binding of the pheromone to the receptor, a G α subunit is released. This has a positive role in signal transduction within the *Sz. pombe* cell, as indeed is the case in many mammalian cells. This contrasts with *S. cerevisiae*, in which the G α subunit is a negative regulator. Accordingly, the *Sz. pombe* system can be thought to be more closely analogous to GPCRs in higher eukaryotes such as mammals.

It has now been realised that an appropriately modified *Sz. pombe* would be a good model for studying GPCRs for identifying components of GPCR pathways, for identifying mutants in the GPCR pathways and for identifying compounds which stimulate or inhibit GPCR-regulated signalling pathways.

Sz. pombe also appears to have greater cell wall permeability than *S. cerevisiae*. This may prove to be invaluable in the study of receptors with large or complex ligands.

A first aspect of the invention provides a *Schizosaccharomyces pombe* (*Sz. pombe*) yeast cell comprising:

- (i) a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway which is derepressed during the mitotic phase of cell growth;

(ii) a reporter system for reporting a signal mediated by the GPCR-regulated signalling pathway;

and wherein:

(a) the GPCR is heterologous, and/or

(b) the reporter system comprises a reporter gene and a promoter, the reporter gene being operatively linked to the promoter, and the promoter being regulatable by the GPCR, at least one of the reporter genes and the promoter being heterologous.

Expression of some of the components of the GPCR-regulated signalling machinery is normally repressed in *Sz. pombe* during mitotic growth and it is necessary to remove this repression in order to study signalling during the mitotic phase of cell growth. Methods for derepressing the GPCR-regulated signalling machinery are discussed below.

It has been discovered that maintaining a derepressed GPCR-regulated signalling pathway during the mitotic phase of cell growth allows *Sz. pombe* to be used to study GPCRs. That is, the cell has one or more signalling components activated during the mitotic phase to enable, for example, the binding of a suitable ligand to the GPCR to increase or reduce transcription of a reporter gene.

The yeast cell will generally comprise one or more mutations to derepress the GPCR-regulated pathway in mitotic growth. A nutritional control pathway can be disrupted by mutation for this purpose.

It is normally necessary to starve *Sz. pombe* cells to induce them to mate and derepress the GPCR-regulated signalling pathway. The relatively high level of cytoplasmic cAMP that exists during mitotic growth is reduced as nutrients become limiting and this helps to trigger sexual development. Strains lacking adenylate cyclase (which converts ATP to cAMP) have no cytoplasmic cAMP but grow reasonably well. They are derepressed for sexual development and respond to mating pheromones during mitotic growth. Accordingly, preferably the yeast cell is adenylate cyclase deficient. More especially, the *cyr1* gene, which encodes adenylate cyclase, is physically or functionally removed or disrupted, for example by insertion of a DNA sequence. The inserted DNA sequence may

be anything convenient, but in a preferred embodiment of the invention the inserted DNA may comprise a reporter gene; the *ura4* gene is one example, as will be discussed below. The *cyr1* gene is discussed in detail in the article by John Davey and Olaf Nielsen (Davey and Nielsen, 1994). Other methods for bypassing the nutritional control of the signalling machinery are available and may be used to derepress the GPCR-regulated signalling pathway in cells of the invention. This could include mutation of any gene that has the effect of repressing sexual differentiation. Such genes include those encoding certain protein kinases repressing sexual differentiation, including the *pat1* gene. The use of a mutation in this gene to bypass the nutritional control and derepress the GPCR-regulated signalling pathway has been described (Davey and Nielsen, 1994).

As indicated above, Davey and Nielsen, 1994 discloses the identification of mutants involved in sexual differentiation and pheromone response. A temperature-sensitive *pat1* mutant (*pat1-114*) allows the arrest of mitotic growth in response to M-factor. A mutation in the adenylate cyclase gene (*cyr1*) was also studied. The authors indicated that cells containing such a mutation have a problem in that they become insensitive or adapted to the pheromone. The perceived problems identified by the authors of the paper, have, in contrast with the analogous situation in *Saccharomyces cerevisiae*, now been found not to present a practical difficulty in heterologous *Sz. pombe* systems: specifically, it has been found that *Sz. pombe* cells containing a mutant *cyr1* and a heterologous GPCR or a suitable heterologous reporter gene do not have a serious problem with desensitisation.

The reporter system allows signal transduction to be measured in a variety of ways. For example, suitable reporter system includes the association or dissociation of signalling components (to include, for example, the association of proteins with stimulated GPCRs, the dissociation of $G\alpha$ subunits from negative regulators such as the $G\beta\gamma$ subunits), the generation of second messengers (such as Ca^{2+} mobilisation, changes in cyclic AMP levels, GTP hydrolysis, phospholipid hydrolysis), the modification of signalling components (such as the phosphorylation of e.g. MAP kinases, MAP kinase kinases or MAP kinase kinase kinases) or altered transcription of a gene. Transcription of a gene can be measured directly (for example, mRNA expression may be detected by Northern blots) or indirectly

(for example, the protein product may be measured by a characteristic stain or intrinsic activity).

Preferably the yeast comprises a nucleic acid molecule encoding a heterologous reporter gene, or an endogenous reporter gene, operatively linked to a promoter that is regulated by a GPCR-regulated signalling pathway. By operatively linked, we mean that the heterologous reporter gene, or the endogenous reporter gene, is linked to the promoter in such a way that the promoter is capable of directing transcription of the reporter gene.

The reporter gene may be any nucleic acid sequence encoding a detectable gene product. The gene product may be an untranslated RNA product such as mRNA or antisense RNA. Such untranslated RNA may be detected by techniques known in the art, such as PCR, Northern or Southern blots. Alternatively, the reporter gene may encode a polypeptide, such as protein or peptide, product. A polypeptide may be detected immunologically or by means of its biological activity. The reporter gene may be any known in the art. The reporter gene need not be a natural gene, and the term "gene" in this sense should not be taken to imply identity with any natural gene. Reporter genes useful in the invention may be the same as certain natural genes, but may differ from them either in terms of non-coding sequences (for example one or more naturally occurring introns may be absent) or in terms of coding sequences.

The reporter gene may encode a protein that allows the yeast cell to be selected by, for example, a nutritional requirement. For example, the reporter gene may be the *ura4* gene which encodes orotidine-5'-phosphate decarboxylase. The *ura4* gene encodes an enzyme involved in the biosynthesis of uracil and offers both positive and negative selection. Only cells expressing *ura4* are able to grow in the absence of uracil, where the appropriate yeast strain is used. Cells expressing *ura4* die in the presence of 5-fluoro-orotic acid (FOA) as the *ura4* gene product converts FOA into a toxic product. Cells not expressing *ura4* can be maintained by adding uracil to the medium. The sensitivity of the selection process can be adjusted by using medium containing 6-azauracil, a competitive inhibitor of the *ura4* gene product. The *his3* gene (encodes imidazoleglycerol-phosphate dehydratase) is also suitable

for use as a reporter gene that allows nutritional selection. Only cells expressing *his3* are able to grow in the absence of histidine, where the appropriate yeast strain is used.

The reporter gene may encode for a protein that allows the yeast to be used in a chromogenic assay. For example, the reporter may be the *lacZ* gene from *Escherichia coli*. This encodes the β -galactosidase enzyme. This catalyses the hydrolysis of β -galactoside sugars such as lactose. The enzymatic activity of the enzyme may be assayed with various specialised substrates, for example X-gal (5-bromo-4-chloro-3-indoyl- β -D-galactoside) or *o*-nitrophenyl- β -D-galactopyranoside, which allow reporter enzyme activity to be assayed using a spectrophotometer, fluorometer or a luminometer.

The gene encoding green fluorescent protein (GFP), which is known in the art, may also be used as a reporter gene.

The reporter gene may also encode a protein that is capable of inducing the cell, or an extract of a cell, to produce light. For example, the reporter gene may encode luciferase. The luciferase reporter genes are known in the art. They are usually derived from firefly (*Photinus pyralis*) or sea pansy (*Renilla reniformis*). The luciferase enzyme catalyses a reaction using D-luciferin and ATP in the presence of oxygen and Mg^{2+} resulting in light emission. The luciferase reaction is quantitated using a luminometer that measures light output. The assay may also include coenzyme A in the reaction that provides a longer, sustained light reaction with greater sensitivity.

An alternative form of enzyme that allows the production of light is aequorin, which is known in the art.

Most preferably, the reporter gene encodes β -lactamase. This reporter gene has certain advantages over, for example, *lacZ*. There is no background activity in mammalian cells or yeast cells, it is compact (29 kDa), it functions as a monomer (in comparison with *lacZ* which is a tetramer), and has good enzyme activity. This may use CCF2/AM, a FRET-based membrane permeable, intracellularly trapped fluorescent substrate. CCF2/AM has a 7-hydroxycoumarin linked to a fluorescein by a cephalosporin core. In the

intact molecules, excitation of the coumarin results in efficient FRET to the fluorescein, resulting in green fluorescent. Cleavage of the CCF2 by β -lactamase results in spatial separation of the two dyes, disrupting FRET and causing cells to change from green to blue when viewed using a fluorescent microscope. The retention of the cleaved product allows the blue colour to develop over time, giving a low detection limit of, for example, 50 enzyme molecules per cell. This results in the reporter gene being able to be assayed with high sensitivity. It also allows the ability to confirm results by visual inspection of the cells or the samples.

The nucleic acid molecule comprising the reporter gene under the control of the GPCR-regulated promoter may additionally comprise one or more additional regulatory elements, such as upstream activating sequences (UAS), termination sequences and/or secretory sequences known in the art. The secretory sequences may be used to ensure that the product of the reporter gene is secreted out of the yeast cell.

Preferably the promoter is regulatable by a yeast mating pheromone binding to its GPCR. The yeast mating pheromone may especially be P-factor pheromone. This is especially preferred because the P-factor pheromone is relatively easy to produce.

The promoter is preferably an endogenous *Sz. pombe* promoter which is regulated by the GPCR. However, it does not have to be endogenous. Certain heterologous promoters may be found to be so regulatable, or may be engineered to be, for example by inclusion of a TR-box motif as described by Aono, *et al.* (1994).

More preferably, the promoter is the *sxa2* promoter, or a homologue or analogue thereof. By homologue or analogue we mean a promoter which may contain one or more changes to the nucleic acid sequence encoding the *sxa2* promoter but which maintains the same functional activity as the *sxa2* promoter. The *sxa2* gene to which the *sxa2* promoter is attached in wild-type cells, encodes a carboxypeptidase that, in wild-type cells, inactivates P-factor by removal of the C-terminal leucine residue (Ladds *et al.*, 1996). Use of the *sxa2* promoter for construction of a GPCR-regulated reporter is advantageous because the promoter is tightly regulated by the P-factor receptor (the GPCR) to which the P-factor

pheromone binds. Only one copy of the *sxa2* promoter exists in wild-type cells. Accordingly, it is possible to remove the naturally occurring *sxa2* promoter and its associated *sxa2* gene and replace it with a construct containing the reporter gene under the transcriptional control of the *sxa2* promoter. This promoter-reporter construct may be integrated into the chromosome of the yeast cell.

Integrating the promoter-reporter gene construct into the chromosome of the yeast cell is advantageous because a known number of reporter genes are then found within each cell. If the promoter-reporter gene construct is placed on a plasmid, then the number of reporter genes in each cell may vary since the copy number of the plasmid may vary considerably and is not constant.

Inactivating the endogenous *sxa2* gene, for example by at least partially deleting the *sxa2* gene, can improve the sensitivity of the assay when P-factor is used to stimulate the GPCR. This is because inactivation of the carboxypeptidase reduces inactivation of the P-factor which may be used to stimulate the GPCR. The reporter gene may be linked to any remaining *sxa2* gene, for example to form a fusion protein. Alternatively, the entire *sxa2* gene may be deleted and the reporter gene inserted in its place.

Preferably, the yeast cell used exhibits a stable mating type. Mating type in *Sz. pombe* is determined by information carried at the *mat1* locus. Haploid cells containing the *mat1-P* segment, which contains the *mat1-Pc* and *mat1-Pm* genes, are '+' (P or plus), and those with *mat1-M*, encoding *mat1-Mc* and *mat1-Mm* are '-' (M or minus). Expression of *mat1-Pc* and *mat1-Mc* are required for expressing the genes that encode the pheromones and their receptors and hence establish the pheromone communication system. All 4 *mat1* genes are required for meiosis. There are two further mating loci, *mat2* and *mat3* where P and M information is stored but not expressed. In wild-type homothallic strains the information at *mat2* and *mat3* is frequently transferred to the *mat1* locus and cells switch mating type approximately once every three generations. Cultures of such strains are therefore normally a mixture of both mating types (P and M). Even normal heterothallic strains are relatively unstable. Strains with a stable mating type can be generated by either

deleting the *mat2* and *mat3* loci or by mutating the switching machinery, to produce a yeast cell exhibiting a stable mating phenotype (Davey, 1998).

Continued exposure to stimulus will lead to desensitisation of the signalling pathway. Several mechanisms are known to contribute to the desensitisation process. Selected mutations in the genes encoding proteins involved in desensitisation can lead to hypersensitivity and an inability to adapt to stimulation. This could be an advantage when using the strains in high throughput screens.

The yeast cell may be *rgs1* deficient. Strains lacking *rgs1* or having reduced Rgs1 (the product of the *rgs1* gene) activity are hypersensitive to pheromone stimulation (Watson, *et al.*, 1999).

The yeast cell may also be *pmp1* deficient. The *pmp1* gene encodes a dual specificity phosphatase that dephosphorylates the MAPK. Strains lacking this phosphatase exhibit an increased response following stimulation of the cells with a ligand for the GPCR.

The GPCR may be a naturally occurring yeast pheromone receptor. Alternatively, the receptor may be replaced, or contain in addition thereto, an heterologous receptor from another cell. When the GPCR is heterologous, it may be from any species other than *Sz. pombe*. The GPCR may be from a plant species or an animal species, particularly mammals, including economically significant non-human mammals. In one of the most important aspects of the invention, however, it will be a human GPCR. The GPCR may be any GPCR which it is desired to investigate by means of the invention. For example, the yeast cell may express an orphan receptor. That is, a receptor of unknown specific activity, but which has been identified by its homology to other GPCR receptors. The yeast cell may be modified to produce such orphan receptors using techniques known in the art. For example, a plasmid containing a nucleic acid sequence encoding for the orphan receptor operably linked to suitable promoter and regulatory sequences may be inserted into the yeast cell. The receptors may be modified to include a signal sequence that functions in *Sz. pombe*. Suitable signal sequences include those of Mam2, Map3 and of other gene products secreted by the *Sz. pombe* cells. If the wild-type heterologous GPCR cannot be

made functional in *Sz. pombe*, it may be mutated for this purpose. In addition, the *Sz. pombe* cells may express endogenous GPCRs in a functional form.

The *Sz. pombe* cell must contain a G protein that is activated by the GPCR and can interact with the rest of the yeast intracellular signalling machinery. The endogenous *Sz. pombe* G α subunit (Gpa1) may be able to couple the heterologous receptor to the intracellular signalling machinery. However, it may be necessary to engineer the *Sz. pombe* cell to produce a heterologous or chimeric G protein subunit (or subunits). At least 16 G α subunits have been identified in mammals and a given GPCR usually activates only one or a small subset of G α subunits. The amino- and carboxy-termini of G α subunits do not share significant homology, but there are several generalisations that can be made. For example, the amino-termini have been implicated in association with G $\beta\gamma$ subunits and with membranes through N-terminal myristoylation. Interaction with the receptor is thought to involve the carboxy-termini as mutants lacking the 5 C-terminal residues of the G α subunit fail to couple to their receptors (see, for example, Hirsch *et al.*, 1991) and peptides based on the C-terminal region of the G α subunit bind to receptors (Hamm *et al.*, 1988; Palm *et al.*, 1990; Rasenick *et al.*, 1994). Work with chimeric G α subunits further supports a critical role for the C-terminal residues in conferring receptor specificity (Voyno-Yasenetskaya *et al.*, 1994; Liu *et al.*, 1995). Thus, the A1 adenosine receptor naturally couples through Gi but can couple via a G α chimera in which the C-terminal 4 residues of Gq were exchanged for those of Gi2 (Conklin *et al.*, 1993) and the SST3 somatostatin receptor does not couple through Gs but can be coupled to adenylate cyclase by replacing the last 5 residues of Gs with those from Gi2 (Komatsuzaki *et al.*, 1997).

Several reports have demonstrated that heterologously expressed GPCRs can couple to the intracellular signalling machinery in *S. cerevisiae*. Some of these receptors can interact with the endogenous G α subunit (encoded by the *GPA1* gene), including those for rat somatostatin (Price *et al.*, 1995), rat A_{2A} adenosine (Price *et al.*, 1996), human lysophosphatidic acid (Erickson *et al.*, 1998) and human UDP-glucose (Chambers *et al.*, 2000). Several other receptors, including that for human growth hormone releasing hormone, do not couple to the *S. cerevisiae* Gpa1 (Kajkowski *et al.*, 1997). In order to attain coupling of these receptors to the intracellular signalling machinery, the *S. cerevisiae*

G α subunit can be replaced by a mammalian G α subunit or by a chimeric G α subunit in which the C-terminal region of the yeast G α subunit is replaced with the equivalent region of the mammalian G α subunit. Many examples of the use of chimeric G α subunits are available (Price *et al.*, 1995; Bass *et al.*, 1996; Kajkowski *et al.*, 1997; Klein *et al.*, 1998; Baranski *et al.*, 1999; Swift *et al.*, 2000). In some instances, production of the chimeric G α may involve the replacement of as few as 5 residues from the C-terminus of the endogenous yeast G α subunit with the equivalent residues from the mammalian G α subunit. Such constructs are sometimes referred to as 'G α -transplants'. There are several reports describing the use of G α -transplants in *S. cerevisiae* (Olesnický *et al.*, 1999; Brown *et al.*, 2000; Chambers *et al.*, 2000; Erlenbach *et al.*, 2001).

The use of G α -transplants based on the endogenous *Sz. pombe* G α subunit may be used to improve the coupling of heterologous GPCRs. To ensure the correct stoichiometric relationship between the G α -transplant and the G $\beta\gamma$ subunits, it may be necessary to replace the chromosomal copy of the natural *Sz. pombe* G α gene (*gpa1*) with the equivalent construct encoding the G α -transplant. However, it is also likely that expression from other promoters is compatible with coupling of the G α subunits to the receptors.

Yeast cells of the invention containing the G α -transplants, and vectors, such as plasmids, cosmids, etc. containing nucleic acid encoding the transplants are included in the invention.

The yeast cell may additionally comprise one or more nucleic acid molecules, such as plasmids, encoding for one or more peptides or proteins, to allow the peptide or protein to be assayed for its effect on GPCR-regulated activity of the reporter system. Alternatively, one or more other chemical compounds may be added to determine the effect of the compound on reporter system activity.

Preferably, the yeast cells contain an auxotrophic marker that allows the selection of plasmids in the yeast cells. The *leu1* mutation provides one such marker and makes growth of the cells dependent upon the addition of leucine or on the introduction of a plasmid containing the *leu1* gene. Similar mutations can also be made to genes involved in the biogenesis of other nutrients (including histidine, lysine and arginine). Such markers include *ade1*, *ade6*, *arg3*, *CAN1*, *his3*, *his7* and *ura4*, all of which are known in the art.

Plasmids containing the nucleic acid encoding for a peptide or protein to be assayed may contain one or more promoter, termination and processing signal sequences. Suitable promoters include the thiamine repressed *nmt1* promoter. This is repressed by the presence of thiamine. Other suitable promoters include *adh1* and *fbp1*, which are known in the art.

The plasmid may also contain a yeast autonomous replication sequence (ARS) to enable the plasmid to replicate in the *Sz. pombe* cells.

A bacterial origin of replication (*ori*), together with one or more bacterial selection markers, such as the ampicillin or tetracycline-resistant genes, may also be included to allow the plasmid to be replicated within bacterial systems prior to insertion into yeast cells. Additionally, the plasmids may include one or more restriction endonuclease sites to enable nucleic acid sequences encoding the peptide or proteins of interest to be inserted. Most preferably, the nucleic acid sequence encoding the peptides or proteins is random and/or may be in the form of a conformational library. Such libraries are known in the art. This allows the production of random peptides to identify peptide regulators of interest. This also allows a library of yeast cells containing different peptides to be produced.

One or more nucleic acid sequences encoding for known peptides or proteins may be introduced into the cell. This allows, for example, a mammalian GPCR-regulated pathway to be reconstituted within a yeast cell.

The strain may additionally contain an *ade6* mutation that helps to make diploid strains of *Sz. pombe* more stable. This is useful where diploid strains of yeast are desirable.

It is not intended that the modifications to *Sz. pombe* described above necessarily be the only modifications made to the cell. Further modifications can be made as required for tailoring the system to particular circumstances.

A further aspect of the invention provides an isolated nucleic acid molecule comprising a promoter regulatable by G-Protein Coupled Receptor (GPCR)-regulated signalling pathway

in *Schizosaccharomyces pombe*, operatively linked to a reporter gene. It is preferred that the promoter be an *sxa2* promoter or homologue or analogue thereof operatively linked to a reporter gene. The *sxa2* promoter and/or reporter genes may be as previously described.

For the avoidance of doubt, in this context, by reporter gene we mean any detectable gene which is not a naturally occurring *sxa2* gene.

A further aspect of the invention provides the use of a *Schizosaccharomyces pombe* (*Sz. pombe*) yeast cell comprising:

- (i) a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway which is derepressed during the mitotic phase of cell growth;
- (ii) a reporter system for reporting a signal mediated by the GPCR-regulated signalling pathway; or
- (iii) an isolated nucleic acid molecule as defined above to study GPCR-regulated activity.

A further aspect of the invention provides an assay comprising the use of a *Schizosaccharomyces pombe* (*Sz. pombe*) yeast cell comprising:

- (i) a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway which is derepressed during the mitotic phase of cell growth;
- (ii) a reporter system for reporting a signal mediated by the GPCR-regulated signalling pathway; or
- (iii) isolated DNA molecule as defined above.

The invention also provides a method of determining the effect of a compound on GPCR-regulated activity comprised in the steps of:

- (i) providing a *Schizosaccharomyces pombe* (*Sz. pombe*) yeast cell comprising:
 - (a) a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway which is derepressed during the mitotic phase of cell growth;
 - (b) a reporter system for reporting a signal mediated by the GPCR-regulated signalling pathway;
- (ii) introducing the compound to the yeast cell; and

- (iii) noting the output of the reporter system, for example by determining an amount of reporter gene product produced by the yeast cell.

The amount of reporter gene product or other reporter system output may be compared with a control yeast without the compound.

The invention also relates to the use of a *Schizosaccharomyces pombe* (*Sz. pombe*) yeast cell comprising:

- (a) a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway which is derepressed during the mitotic phase of cell growth;
- (b) a reporter system for reporting a signal mediated by the GPCR-regulated signalling pathway;

to identify a compound which acts as the receptor. The compound may be the or a natural ligand for the receptor or be an agonist or antagonist (or partial agonist or partial antagonist). Such compounds affect the ability of the receptor to regulate the GPCR-regulated signalling pathway. The invention therefore encompasses the use of such a yeast cell containing an orphan GPCR to identify compounds that affect the ability of the orphan receptor to regulate the promoter is also provided.

The yeast cell, as defined above, may be used to identify a regulator or a mutant of a GPCR-regulated pathway.

The invention also provides a method of identifying a reagent that modulates GPCR-regulated signalling, comprising:

- (i) providing a *Schizosaccharomyces pombe* (*Sz. pombe*) yeast cell comprising:
 - (a) a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway which is derepressed during the mitotic phase of cell growth;
 - (b) a reporter system for reporting a signal mediated by the GPCR-regulated signalling pathway;
- (ii) producing a random peptide within the yeast cell; and
- (iii) noting the output of the reporter system, for example by measuring an amount of reporter gene product produced by the yeast cell.

A still further aspect of the invention provides a compound capable of modulating GPCR-regulated activity identified by a method according to the invention. Assay kits comprising a yeast cell or isolated nucleic acid molecule as defined above are also provided.

M-cells do not normally express the P-factor mating pheromone (encoded by the *map2* gene). P-factor is an unmodified peptide of 23 amino acids that is initially synthesised as a precursor containing an N-terminal signal sequence and four tandem copies of the mature pheromone. The signal sequence is lost after targeting the precursor into the secretory pathway and the precursor is then processed into the individual subunits before being released into the medium. A plasmid-based *map2* construct that contains a single copy of the pheromone peptide and is expressed under the control of the *nmt1* promoter has been prepared. Reporter strains containing the plasmid secrete P-factor when grown in thiamine-free medium and this elicits an autocrine response in the yeast cell in which the P-factor produced by the cell stimulates the pheromone receptor expressed in the same cell.

A further aspect of the invention therefore provides a yeast cell containing such a construct. Restriction sites may be provided within the construct to allow the P-factor sequence to be replaced by an alternative peptide sequence that is then secreted into the medium. Introducing random sequences into this construct produces a library of yeast strains in which each individual releases a different peptide, and allows random peptides to be assayed for their ability to act as autocrine inducers.

Strains of cells of, and useful in, the invention may be termed "reporter strains".

Another feature of the invention is that it provides a method of determining whether a GPCR is coupled to the intracellular signalling machinery even in the absence of a ligand. Such a method is particularly useful for investigating orphan GPCRs, for which the natural ligand may not be known. The method is based on the as yet unexplained observation that the ligand-independent reporter system response is higher in a cell lacking a coupled receptor than it is in a comparable cell having a coupled receptor.

According to this aspect of the invention, there is therefore provided a method of determining whether a G-Protein Coupled Receptor (GPCR) is coupled to a cell signalling pathway, the method comprising comparing the ligand-independent reporter system output of a *Schizosaccharomyces pombe* (*Sz. pombe*) yeast cell comprising:

- (i) a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway which is derepressed during the mitotic phase of cell growth;
- (ii) a reporter system for reporting a signal mediated by the GPCR-regulated signalling pathway;

with the reporter system output of a reference cell which lacks a functional GPCR.

The reference cell, which itself forms another aspect of the invention, will generally be a *Schizosaccharomyces pombe* (*Sz. pombe*) yeast cell comprising:

- (i) a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway which is derepressed during the mitotic phase of cell growth, wherein the GPCR is absent or otherwise rendered non-functional;
- (ii) a reporter system for reporting a signal mediated by the GPCR-regulated signalling pathway.

The reporter system will be expected to give an output indicative of higher activity from the reference cell than from the cell under investigation if the GPCR in the cell under investigation is coupled to the signalling pathway.

Preferred features of each aspect of the invention are as for each other aspect, *mutatis mutandis*.

The invention will now be described by way of example only, with reference to the following figures:

Figure 1. Schematic diagram showing the identification and step-wise replacement of the *sxa2* gene with *ura4⁺*, and the *sxa2>ura4* and *sxa2>lacZ* reporter genes.

Figure 2. Southern Blot of a *Pvu*II and *Hind*III digest of the constructs shown schematically in Figure 1.

Figure 3. Schematic diagram of the arrangement of the *map2* gene product.

Figure 4. Amino acid sequence of the *map2* gene product.

Figure 5. Schematic diagram showing the construction of a construct containing only one copy of the P-factor gene (the mono P construct).

Figure 6. Amino acid sequence of the mono P construct.

Figure 7A. Schematic diagram showing the replacement of a P-factor gene with a nucleic acid sequence encoding a random peptide, where "n" is an unknown amino acid.

Figure 7B. Amino acid sequence of the modified P-factor gene product encoding a random peptide, where "n" is an unknown amino acid.

Figure 8. Positive and negative selection using the *ura4* reporter gene: a) Growth of yeast cells on plates without uracil upon stimulation with P-factor; b) Inhibition of growth on FOA plates. Yeast cells are stimulated with 1, 10 and 100 units of P-factor.

Figure 9. Growth of *sxa2>ura4* yeast cells on plates without uracil. The yeast cells are stimulated with between 0.1 and 1000 units/ml. P-factor.

Figure 10. Identification of mutants having enhanced sensitivity to P-factor stimulation. *sxa2>ura4* cells were grown on plates lacking uracil.

Figure 11. Identification and characterisation of *rgs1* mutants using the *sxa2>ura4* strain.

Figure 12. Thiamine-inducible expression of P-factor using the *sxa2>ura4* reporter strain and a thiamine-inducible P-factor construct.

Figure 13. P-factor stimulation of β -galactosidase in the *sxa2>lacZ* reporter strain.

Figure 14. Coupling of the human CRH receptor in *Sz. pombe* strains containing various G α -transplants.

Figure 15. Demonstrating the coupling of a receptor in the absence of its ligand.

METHODS

All manipulations were by standard methods (see, for a general review, Davey *et al.*, 1995). Reagents were obtained from common laboratory suppliers and used according to the manufacturer's recommendations. Unless stated otherwise, the polymerase chain reaction (PCR) was performed using *Pwo* DNA polymerase (from *Pyrococcus woesei*; supplied by Boehringer Mannheim) as this has a 3'-5' exonuclease (proof-reading) activity and reduced the introduction of errors during amplification. *TAQ* polymerase (from *Thermus aquaticus*; supplied by Boehringer Mannheim) was used for PCR with primers containing random sequences.

The yeast strains identified below are merely examples. Other suitable strains can be readily identified or produced using techniques known in the art.

Yeast strains

JY271 is *h⁻, cyl1::ura4, ade6-M216, leu1-32, ura4-D18* and is equivalent to JZ300 (Maeda *et al.*, 1990). This is an M-cell but not stable and can switch mating type. The *cyl1* gene (encoding adenylate cyclase) was disrupted by insertion of the *ura4* gene (pDAC5), resulting in a cell which requires adenine and leucine for growth.

JY330 is *mat1-P, Δmat2/3::LEU2⁻, leu1-32*. The *mat2-P* and *mat3-M* donor mating cassettes were deleted by insertion of *LEU2* (Klar and Miglio, 1986) and a *LEU2⁻* isolate was then identified (Klar and Bonaduce, 1991).

JY444 is *mat1-M, Δmat2/3::LEU2⁻, leu1-32, ura4-D18* and is a stable M-cell that requires leucine and uracil for growth.

The *ura4* cassette used to disrupt the *cyl1* gene in JY271 was removed by standard techniques to create JY271B. JY271B is *h⁻, cyl1-D51, ade6-M216, leu1-32, ura4-D18*.

This is an M-cell but not stable and can switch mating type. The *cyr1* gene (adenylate cyclase) is disrupted. The cell requires adenine, leucine and uracil for growth.

JY271B was crossed with JY330 to generate JY361. JY361 is *mat1-P*, Δ *mat2/3::LEU2*⁻, *leu1-32*, *ade6-M216*, *ura4-D18*, *cyr1-D51*. This is a stable P-cell in which the *cyr1* gene (adenylate cyclase) is disrupted. The cell requires adenine, leucine and uracil for growth.

JY361 was crossed with JY444 to generate JY486. JY486 is *mat1-M*, Δ *mat2/3::LEU2*⁻, *leu1-32*, *ade6-M216*, *ura4-D18*, *cyr1-D51*. This is a stable M-cell in which the *cyr1* gene (adenylate cyclase) is disrupted. The strain requires adenine, leucine and uracil for growth.

The *sxa2* gene in JY486 was disrupted using a *ura4*⁺ cassette to generate JY522. The manipulation of the *sxa2* gene is described in more detail below. The disruption cassette was a *NcoI*-to-*BamHI* fragment from JD883. JY522 is *mat1-M*, Δ *mat2/3::LEU2*⁻, *leu1-32*, *ade6-M216*, *ura4-D18*, *cyr1-D51*, *sxa2::ura4*⁺. This is a stable M-cell in which the *cyr1* gene (adenylate cyclase) is disrupted. The *sxa2* gene (encodes a serine carboxypeptidase) is also disrupted. The strain requires adenine and leucine for growth.

The disrupted *sxa2* gene in JY522 was replaced with the *sxa2>lacZ* reporter to generate JY546. The *sxa2>lacZ* reporter construct is from JD954. JY546 is *mat1-M*, Δ *mat2/3::LEU2*⁻, *leu1-32*, *ade6-M216*, *ura4-D18*, *cyr1-D51*, *sxa2>lacZ*. This is a stable M-cell in which the *cyr1* gene (adenylate cyclase) is disrupted. The strain has an *sxa2>lacZ* reporter integrated at the *sxa2* locus and expresses *lacZ* in response to pheromone stimulation. This strain requires adenine, leucine and uracil for growth.

The disrupted *sxa2* gene in JY522 was also replaced with the *sxa2>ura4* reporter to generate JY603. The *sxa2>ura4* reporter construct is from JD929. JY603 is *mat1-M*, Δ *mat2/3::LEU2*⁻, *leu1-32*, *ade6-M216*, *ura4-D18*, *cyr1-D51*, *sxa2>ura4*, and is a stable M-cell. The *cyr1* gene (adenylate cyclase) is disrupted. This has an *sxa2>ura4* reporter integrated at the *sxa2* locus and expresses *ura4* in response to pheromone stimulation. The strain requires adenine and leucine for growth.

Constructing the *sxa2*>reporter strains

Figures 1 and 2 summarise the methods used to manipulate the *sxa2* gene and promoter.

The *sxa2* ORF was first replaced with a 1.8 kb *Sz. pombe ura4⁺* cassette (Grimm *et al.*, 1988). The complete *sxa2* locus was amplified by PCR using the sense primer JO760 (ggggggtacCATGGCTAGAAATCCGCCATTGTGTG; lower-case letters are not complementary to *sxa2* but the oligonucleotide includes a *KpnI* site [ggtac*C] and an *NcoI* site [c*CATGG] where digestion leaves ends that are fully homologous to the chromosomal sequence) and the antisense primer JO683 (CTTCTCGTAAAGGCACATTGACGG, complementary to a region immediately downstream of the *BamHI* site at position 2043). The resulting PCR product was cloned into the *KpnI* and *BamHI* sites of pSP72 (Promega) to generate JD808 (pSP72 containing the *sxa2* locus SEQ ID 31). This was used as template for PCR with JO746 (TGAAAAGAGAGACAATG; antisense primer complementary to a region immediately upstream of the ATG initiator codon for *sxa2*) and JO745 (TAAAAGTTTAATATC; sense primer complementary to a region that includes the TAA stop codon for *sxa2*) and the product ligated to the *ura4⁺* cassette (to generate JD857, pSP72 containing a construct suitable for disruption of *sxa2* SEQ ID 32) or to PCR products corresponding to either the *lacZ* ORF (to generate JD954, pSP72 containing the *sxa2*>*lacZ* reporter construct SEQ ID 33) or the *ura4* ORF (to generate JD929, pSP72 containing the *sxa2*>*ura4* reporter construct SEQ ID 34). The *lacZ* ORF was prepared by amplification using the sense primer JO660 (ATGCAGCTGGCACGACAGGTTTCCCGAC; includes the ATG initiator codon and next 25 bases of the *lacZ* ORF) and the antisense primer JO661 (TTTTTGACACCAGACCAACTGGTAATGGTAGC; complementary to the 3' end of the *lacZ* ORF but lacks the stop anticodon). The *ura4* ORF was prepared by amplification using the sense primer JO828 (ATGGATGCTAGAGTATTTC; includes the ATG initiator codon and next 16 bases of the *ura4* ORF) and the antisense primer JO759 (ATGCTGAGAAAGTCTTTGC; complementary to the 3' end of the *ura4* ORF but lacks the stop anticodon).

JY486 (a mating stable M-cell lacking *cyr1*) was transformed with a *NcoI*-*Bam*HI fragment corresponding to the *sxa2::ura4⁺* construct (isolated from JD857), and stable Ura4⁺ transformants were initially screened by PCR and replacement of the *sxa2* locus was confirmed by Southern blot (Figure 2). A correct *sxa2::ura4⁺* disruptant (JY522) was then transformed with the *NcoI*-*Bam*HI fragments corresponding to the *sxa2>lacZ* reporter (isolated from JD954) or the *sxa2>ura4* reporter (isolated from JD929). Stable Ura⁺ transformants were selected by their ability to grow in the presence of 5'fluoro-orotic acid (Boeke *et al.*, 1987) and homologous integration of the reporter constructs at the *sxa2* locus was confirmed by Southern blot for JY546 (*sxa2>lacZ*) and JY603 (*sxa2>ura4*). Southern blot analysis was performed on genomic DNA digested with *Pvu*II & *Hind*III and a probe corresponding to the 5' untranslated region of *sxa2*.

Constructing the G α -transplants

This was undertaken using techniques well known in the art. A *Spe*I-*Pst*I fragment from *gpa1* (SEQ ID 1, 15) was cloned into the *Spe*I and *Pst*I sites of the plasmid pKS-Bluescript (Stratagene). This 324 bp fragment contains the last 24 residues of Gpa1 and 250 base pairs from the 3' untranslated region of the *gpa1* gene. The resulting clone (JD1647) was then used as template for a series of polymerase chain reactions using oligonucleotide primers that made the desired changes to the residues at the C-terminus of Gpa1. Each reaction used the antisense primer JO1354 (TAGATTGTTGGACATAATCGTATCTTGAACGG; complementary to a region from position 1206 to position 1175 relative to the initiator ATG of *gpa1*) and an appropriate sense primer that introduced the desired changes and was complementary to the region immediately downstream of the Gpa1 open reading frame; JO1344 for the G α q-transplant (gaatataatcttggtTAGATGAATTTTTCCTTAAC, lower case letters change the last 5 residues of *Sz. pombe* Gpa1 to EYNLV), JO1345 for the G α s-transplant (caatatgaactcttTAGATGAATTTTTCCTTAAC; change last 5 residues of Gpa1 to QYELL), JO1346 for the G α o-transplant (ggatgcggactttatTAGATGAATTTTTCCTTAAC, change last 5 residues of Gpa1 to GCGLY), JO1347 for the G α i2-transplant (gattgcggacttttTAGATGAATTTTTCCTTAAC, change last 5 residues of Gpa1 to DCGLF), JO1348 for the G α i3-transplant

(gaatgcggactttatTAGATGAATTTTTCCTTAAC, change last 5 residues of Gpa1 to ECGLY), JO1349 for the G α z-transplant (tatattggactttgcTAGATGAATTTTTCCTTAAC, change last 5 residues of Gpa1 to YIGLC), JO1350 for the G α 12-transplant (gatattatgcttcaaTAGATGAATTTTTCCTTAAC, change last 5 residues of Gpa1 to DIMLQ), JO1351 for the G α 13-transplant (caacttatgcttcaaTAGATGAATTTTTCCTTAAC, change last 5 residues of Gpa1 to QLMLQ), JO1352 for the G α 14-transplant (gaatttaattcttggTAGATGAATTTTTCCTTAAC, lower case letters change the last 5 residues of Gpa1 to EFNLV) and JO1353 for the G α 16-transplant (gaaattaattcttctTAGATGAATTTTTCCTTAAC, change last 5 residues Gpa1 to EINLL).

The PCR products were sequenced to confirm that the correct changes had been made and were then used to replace the equivalent *SpeI-PstI* fragment from JD1645 (pSP71-Gpa1). JD1645 contains the complete *gpa1* sequence from an *EcoRI* site at position -676 (relative to the initiator ATG) to a *BglIII* site at position 1938. This generated a series of plasmids containing the modified *gpa1* sequences; JD1649 (G α q-transplant SEQ ID 17, 03), JD1650 (G α s-transplant SEQ ID 16, 02), JD1651 (G α o-transplant SEQ ID 18, 04), JD1652 (G α i2-transplant SEQ ID 19, 05), JD1653 (G α i3-transplant SEQ ID 20, 06), JD1654 (G α z-transplant SEQ ID 21, 07), JD1655 (G α 12-transplant SEQ ID 22, 08), JD1656 (G α 13-transplant SEQ ID 23, 09), JD1657 (G α 14-transplant SEQ ID 24, 10) and JD1658 (G α 16-transplant SEQ ID 25, 11). The coding regions for the different G α -transplants were isolated as *EcoRI-BglIII* fragments and used separately to transform the yeast strain JY1170. JY1170 is *mat1-M*, Δ *mat2/3::LEU2*, *leul-32*, *ade6-M216*, *ura4-D18*, *cyr1-D51*, *mam2-D10*, *gpa1::ura4⁺*, *sxa2>lacZ*. This is a derivative of the standard JY546 reporter strain but it lacks the *mam2* gene (encodes the P-factor receptor) and the *gpa1* gene has been disrupted by insertion of a *ura4⁺* cassette. Ura⁻ transformants were selected on fluoro-orotic acid and Southern blot analyses were used to confirm integration of the G α -transplant constructs at the *gpa1* locus. This generated a series of *Sz. pombe* *sxa2>lacZ* reporter strains lacking the *mam2* pheromone receptor but containing integrated G α -transplants; JY1165 (G α q-transplant), JY1157 (G α s-transplant), JY1158 (G α o-transplant), JY1159 (G α i2-transplant), JY1160 (G α i3-transplant), JY1161

(G α z-transplant), JY1162 (G α 12-transplant), JY1163 (G α 13-transplant), JY1164 (G α 14-transplant) and JY1167 (G α 16-transplant).

Generating peptides for autocrine signalling

M-cells do not normally express the P-factor mating pheromone (encoded by the *map2* gene). P-factor is an unmodified peptide of 23 amino acids that is initially synthesised as a precursor containing an N-terminal signal sequence and four tandem copies of the mature pheromone. The signal sequence is lost after targeting the precursor into the secretory pathway and the precursor is then processed into the individual subunits before being released into the medium. A plasmid-based *map2* construct that contains a single copy of the pheromone peptide and is expressed under the control of the thiamine-regulated *nmt1* promoter shown schematically in Figures 3 to 6 was prepared.

This was undertaken using techniques well known in the art. Reporter strains containing the plasmid secrete P-factor when grown in thiamine-free medium (the *nmt1* promoter is on) and this elicits an autocrine response in the strain. Restriction sites within the construct allow the P-factor sequence to be replaced by an alternative peptide sequence that would then be secreted into the medium (Figures 7A and 7B). Introducing random sequences into this construct produces a library of strains in which each individual releases a different peptide. This allows ligands capable of binding to the pheromone receptor or another GPCR to be identified.

Expression and Application of Reporter Gene Constructs

Demonstration of a *sxa2>ura4* Reporter Construct

JY603 yeast cells containing the construct were spread as a confluent layer of cells (about 10^7 cells on each plate) on DMM medium lacking uracil. Paper disks were placed on the dried surface of the cells and aliquots containing different amounts of P-factor were added to each disk. The plates were then incubated at 29°C for 3 days. Figure 8A shows that cells are not normally able to grow in the absence of uracil but the P-factor induces expression of the *sxa2>ura4* reporter and allows a growth of cells around the disks. The halo is largest around the disk containing 100 units of P-factor.

This is also demonstrated in Figure 4, except that a series of plates containing different concentrations of P-factor were used. All of the plates received the same number of yeast cells (about 2,000 cells per plate). The cells are not normally able to grow in the absence of uracil but the P-factor induces expression of the *sxa2>ura4* reporter and allows cells to form colonies. There are no colonies on the plates containing 0.1 or 1.0 units/ml. but colonies form on plates containing P-factor with at least 10 units/ml.

Figure 8B shows plates which contain uracil and 5-fluoro-orotic acid (FOA). Cells not expressing *ura4* are able to grow on these plates but those expressing *ura4* convert the FOA into a toxic compound and die. There are clear halos of no growth around the disk containing the P-factor. The halo is largest around the disk containing 100 units of P-factor.

Identification of Mutants

The *sxa2>ura4* reporter system allows the identification of mutants. Cells can be randomly mutagenised and spread on plates containing P-factor at 0.1 units/ml to identify mutations that make the cells more sensitive to stimulation. Figure 10 shows two of these mutations. This approach can also be used to identify mutant forms of various proteins involved in regulating the signalling pathway as shown in Figure 11.

The *sxa2>ura4* reporting strain was randomly mutagenised and then spread on plates lacking uracil but containing P-factor at 0.1 units/ml. The wild-type cells do not normally grow on these plates, since they require P-factor at a concentration of at least 10 units/ml. A number of mutants that had increased sensitivity to signalling and were now able to grow at a low level of P-factor were identified. Two of these mutants have been characterised as being *rgs1* and *pmp1*. The *pmp1* mutant does not grow on plates lacking P-factor but grows on a very low level of P-factor. This demonstrates that it is a hypersensitive mutant. In contrast, the *rgs1* mutant grows even in the absence of P-factor, showing that it is a constitutive responder (that is, it expresses the reporter gene in the absence of stimulation by ligand).

The Applicants mutated the cloned *rgs1* gene to isolate mutant forms of the protein with altered properties and screened for isolates that were either gain of function mutants (have increased activity relative to the normal Rgs1 protein) or dominant negative mutants (inactive mutants that inhibit the activity of the normal Rgs1 protein in the same cell).

Autocrine Signalling

The inventors modified a version of the *map2* gene that encodes the P-factor precursor was modified so that it contained a single copy of the P-factor ("mono P"). This is cloned into a plasmid so that expression of the P-factor was under the control of the thiamine-repressible *nmt1* promoter. The plasmid was introduced into M-cells which do not normally produce P-factor but are able to respond to P-factor. The cells were spread on plates lacking uracil but containing either no thiamine or 5 μ M thiamine (see Figure 12). The thiamine induces expression and release of the P-factor, causing autocrine signalling of the cell. This results in the expression of the *sxa2>ura4* reporter.

The *lacZ* Reporter

The *sxa2>lacZ* reporter strain was grown in the presence of varying amounts of P-factor. The amount of β -galactosidase released was assayed using

o-nitrophenyl- β -D-galactopyranoside and measuring the amount of product at OD₄₂₀. Figure 13 shows the effect of adding P-factor over time and with increasing concentration. The concentration-dependent assay was measured 16 hours after adding the pheromone.

Coupling of a human GPCR

By way of example, the human receptor for corticotrophin releasing hormone (CRH, also known as corticotrophin releasing factor or CRF) was expressed in the *Sz. pombe* *sxa2>lacZ* reporter strains containing either Gpa1 or the various G α -transplants. The yeast strains were transformed with pREP3X:CRH-R1 (SEQ ID 30), a plasmid that places the CRH receptor (see SEQ ID 28 and SEQ ID 14) type 1 α under the control of the *nmt1* promoter. Transformants were grown in the absence of thiamine (to allow expression of the receptor) and then exposed to CRH at 10⁻⁶ M (control cells were exposed to solvent lacking CRH). The amount of β -galactosidase released was assayed after 16 hours using *o*-nitrophenyl- β -D-galactopyranoside (Figure 14). A low level of coupling was observed with the endogenous Gpa1 but this was considerably improved in the G α s- and G α 16-transplants.

CRH is a 41-residue peptide that is a major regulator of the body's stress axis. Although it has several functions, its best characterised role is in initiating pituitary-adrenal responses to stress, an effect mediated through CRH-R1 α (Vale *et al.*, 1981). This receptor normally functions through G α s, resulting in activation of adenylate cyclase and increased levels of cAMP (Giguere *et al.*, 1982; Bilezikjian and Vale, 1983; Grammatopoulos *et al.*, 1996). The observed coupling to the G α s-transplant is consistent with the activity of the CRH-R1 α receptor in mammalian cells. G α 16 is known to interact with a wide range of GPCRs (Milligan *et al.*, 1996).

There are no reports of the coupling of the human CRH receptor to the signalling machinery in the budding yeast *S. cerevisiae* and a direct comparison with the *Sz. pombe* reporter strains reported here is therefore not possible. It is perhaps significant however that a peptide ligand similar to CRH appears unable to gain access to receptors at the surface of *S. cerevisiae* cells (Baranski *et al.*, 1999). The C5a chemoattractant receptor is

functional in *S. cerevisiae* but can only be stimulated by its ligand (a 74-residue peptide) when both the receptor and the ligand are expressed in the same cell. Such autocrine stimulation is required because the C5a ligand is unable to traverse the *S. cerevisiae* cell wall.

Sz. pombe is also surrounded by a cell wall but it has a very different structure to that surrounding *S. cerevisiae* (for reviews, see, Osumi, 1998; Smits *et al.*, 1999) and previous studies of intoxication by diphtheria toxin demonstrated that the two have quite different permeability properties. Diphtheria toxin, secreted by certain strains of *Corynebacterium diphtheriae*, catalyses the ADP-ribosylation of eukaryotic aminoacyl transferase II (EF-2) using NAD as substrate. This reaction forms the basis for its toxicity toward eukaryotic organisms. Intoxication requires the entry of the toxin into the cytoplasm after internalisation by endocytosis. Studies have investigated the effects of diphtheria toxin on protein synthesis in *S. cerevisiae* (Murakami *et al.*, 1982) and *Sz. pombe* (Davey, 1991). Although the *Sz. pombe* cells were sensitive to the toxin, intact *S. cerevisiae* cells were resistant to its effects. In contrast, *S. cerevisiae* spheroplasts (cells in which the cell wall has been enzymatically removed) were sensitive to the toxin, suggesting that the failure of the toxin to enter intact cells was due to its inability to cross the cell wall. Diphtheria toxin is a heterodimer composed of an N-terminal A fragment (molecular weight 24,000 daltons) that is enzymatically active and a C-terminal B fragment (molecular weight 39,000 daltons) that has no apparent enzymatic activity but is required for toxicity.

The apparent greater permeability of the cell wall in *Sz. pombe* might be invaluable for the study of receptors with large or complex ligands and could provide an advantage over the use of *S. cerevisiae* in such situations.

Demonstrating coupling of a receptor in the absence of its ligand

The lack of available ligands for orphan GPCRs makes it difficult to confirm that such receptors are coupled to the signalling machinery. High throughput screens are therefore performed without the confidence that a ligand which would normally stimulate the receptor would be identified as being active. The inventors have observed an interesting

feature of the *Sz. pombe* reporter strains that can indicate whether a GPCR is coupled to the intracellular signalling machinery even in the absence of its ligand. Such knowledge generates confidence prior to performing high throughput screens. An example of this feature is shown in Figure 15. When a *Sz. pombe* *sxa2>lacZ* reporter strain expressing the normal P-factor pheromone receptor is exposed to P-factor, there is a ligand-dependent induction of β -galactosidase. As expected, a similar strain lacking the P-factor receptor fails to exhibit ligand-dependent induction of the *sxa2>lacZ* reporter. However, the ligand-independent expression of the *sxa2>lacZ* reporter (i.e. the level of β -galactosidase activity observed in the absence of P-factor) is considerably higher in the strain lacking the P-factor receptor than in the strain containing the P-factor receptor. Expressing the human CRH-R1 α receptor in the strain lacking the P-factor receptor reduces the ligand-independent expression of the *sxa2>lacZ* reporter back to the levels observed in the strain containing the P-factor receptor. As there is no P-factor receptor in this strain, addition of P-factor fails to induce further expression of the *sxa2>lacZ* reporter.

This observation is not limited either to the particular reporter system or to the CRH-R1 α receptor and has been observed with many other GPCRs that were then subsequently shown to be coupled to the *Sz. pombe* signalling machinery. A molecular explanation for this effect is not available but it could simply reflect the ability of the receptor to sequester the heterotrimeric G proteins and prevent inappropriate activation of the downstream effector protein(s). Whatever the explanation, the ability of a receptor to reduce the ligand-independent expression of β -galactosidase appears to reflect its ability to couple to the *Sz. pombe* signalling machinery.

Applications

Identify agonists and antagonists for GPCRs, including orphan GPCRs

Reporter strains expressing either a characterised or an orphan receptor can be used in a variety of assays to identify ligands that affect signalling through the receptors. Agonists will elicit a response in the strain while antagonists could be identified by their ability to inhibit stimulation by a ligand known to activate the receptor. Both peptides and small molecules can be screened and assays might be either liquid- or plate-based, depending on the reporter gene used. Screens for peptide ligands could exploit the autocrine signalling of strains producing a library of random peptides.

Identify intracellular signal regulators and modified regulators with altered activities

Regulators of the intracellular response pathway can be identified by their ability to influence signalling in the reporter strains. Over-expression of these proteins will either reduce or increase signalling depending on whether they are positive or negative regulators. A number of mammalian regulators are known to be active in yeast. Regulators identified through these screens can then be mutagenised and the reporter strains used to identify isolates with altered activities. Gain-of-function mutants, for example, would have increased abilities to regulate signalling while dominant-negative mutants would not only be inactive but would also inhibit the activity of the wild type regulator. These mutants could then be introduced back into mammalian systems to assess their ability to regulate other signalling pathways.

Identify reagents that modulate signalling

Random peptides can be expressed in the cytoplasm of the reporter strains and assayed for their ability to regulate signalling. These 'peptamers' could interact directly with components from the signalling pathway or might exert their effect through the intracellular signal regulators mentioned earlier. The screen is not limited to peptide regulators and would also identify small molecules that could influence signalling.

Schizosaccharomyces pombe strain JY546 was deposited under the Budapest Treaty at the National Collection of Yeast Cultures, Norwich, United Kingdom on 27 October 2000. It has been given Accession Number NCYC 2984.

References

- Allshire, R.C., Cranston, G., Gosden, J.R., Maule, J.C., Hastie, N.D. and Fantes, P.A. (1987) A fission yeast chromosome can replicate autonomously in mouse cells. *Cell* **50**, 391-403.
- Aono, T., Yanai, H., Miki, F., Davey, J. and Shimoda, C. (1994) Mating Pheromone-induced Expression of the *mat1-Pm* gene of *Schizosaccharomyces pombe*: Identification of Signalling Components and Characterisation of Upstream Controlling Elements. *Yeast* **10**, 757-770.
- Arkinstall, S., Edgerton, M., Payton, M. and Maundrell, K. (1995) Co-expression of the neurokinin NK2 receptor and G-protein components in the fission yeast *Schizosaccharomyces pombe*. *FEBS Letts.* **375**, 183-187.
- Baldwin, J.M. (1994) Structure and function of receptors coupled to G proteins. *Current Opinion Cell Biol.* **6**, 180-190.
- Baranski, T.J., Herzmark, P., Lichtarge, O., Gerber, B.O., Trueheart, J., meng, E.V., Iiri, T., Sheikh, S.P. and Bourne, H.R. (1999) C5a receptor activation. Genetic identification of critical residues in four transmembrane helices. *J. Biol. Chem.* **274**, 15757-15765.
- Bass, R.T., Buckwalter, B.L., Patel, B.P., Pausch, M.N., Price, L.A., Strnad, J. and Hadcock, J.R. (1996) Identification and characterisation of novel somatostatin antagonists. *Mol. Pharmacol.* **50**, 709-715.
- Bilezikjian, L.M. and Vale, W.W. (1983) Glucocorticoids inhibit corticotropin-releasing factor-induced production of adenosine 3',5'-monophosphate in cultured anterior pituitary cells. *Endocrinology* **113**, 657-662.
- Boeke, J.D., Trueheart, J., Natsoulis, G. and Fink, G.R. (1987) 5'-fluoroorotic acid as a selective agent in yeast molecular genetics. *Methods Enzymol.* **152**, 164-175.
- Brown, A.J., Dyos, S.L., Whiteway, M.S., White, J.H.M., Watson, M-A.E.A., Marzioch, M., Clare, J.J., Cousens, D.J., Paddon, C., Plumpton, C., Romanos, M.A. and Dowell, S.J. (2000) Functional coupling of mammalian receptors to the yeast mating pathway using novel yeast/mammalian G protein α -subunit chimeras. *Yeast* **16**, 11-22.
- Chambers, J.K., Macdonald, L.E., Sarau, H.M., Ames, R.S., Freeman, K., Foley, J.J., Zhu, Y., McLaughlin, M.M., Murdock, P., McMillan, L., Trill, J., Swift, A., Aiyar, N., Taylor, P., Vawter, L., Naheed, S., Szekeres, P., Hervieu, G., Scott, C., Watson, J.M.,

- Murphy, A.J., Duzic, E., Klein, C., Bergsma, D.J., Wilson, S. and Livi, G.P. (2000) A G protein-coupled receptor for UDP-glucose *J. Biol. Chem.* **275**, 10767-10771.
- Conklin, B.R., Farfel, Z., Lustig, K.D., Julius, D. and Bourne, H.R. (1993) Substitution of three amino acids switches receptor specificity of G α_q to G α_i . *Nature* **363**, 274-276.
- Davey, J. (1991) Isolation and quantitation of M-factor, a diffusible mating factor from the fission yeast *Schizosaccharomyces pombe*. *Yeast* **7**, 357-366.
- Davey, J. and Nielsen, O. (1994) Mutations in *cyr1* and *pat1* reveal pheromone-induced G1 arrest in the fission yeast *Schizosaccharomyces pombe*. *Curr. Genet.* **26**, 105-112.
- Davey, J., Egel, R. and Nielsen, O. (1995) Pheromone procedures in fission yeast. in *Microbial Gene Techniques; Methods in Molecular Genetics 6B* (Adolph, K.W., ed.) Academic Press, San Diego. pp 247-263.
- Davey, J. (1998) Fusion of a fission yeast. *Yeast* **14**, 1529-1566.
- Erickson, J.R., Wu, J.J., Goddard, G., Tigyi, G., Kawanishi, K., Tomei, L.D. and Kiefer, M.C. (1998) Edg-2/Vzg-1 couples to the yeast pheromone response pathway selectively in response to lysophosphatidic acid. *J. Biol. Chem.* **273**, 1506-1510.
- Erlenbach, I., Kostenis, E., Schmidt, C., Hamdan, F.F., Pausch, M.H. and Wess, J. (2001) Functional expression of M1, M3 and M5 muscarinic acetylcholine receptors in yeast. *J. Neurochem.* **77**, 1327-1337.
- Ficca, A.G., Testa, L., and Valentini, G.P.T. (1995) The human β_2 -adrenergic receptor expressed in *Schizosaccharomyces pombe* retains its pharmacological properties. *FEBS Letts.* **377**, 140-144.
- Giguere, V., Labrie, F., Cote, J., Coy, D.H., Sueiras-Diaz, J. and Schally, A.V. (1982) Stimulation of cyclic AMP accumulation and corticotropin release by synthetic ovine corticotropin-releasing factor in rat anterior pituitary cells: site of glucocorticoid action. *Proc. Natl. Acad. Sci. USA* **79**, 3466-3469.
- Grammatopoulos, D., Stirrat, G.M., Williams, S.A. and Hillhouse, E.W. (1986) The biological activity of the corticotropin-releasing hormone receptor-adenylate cyclase complex in human myometrium is reduced at the end of pregnancy. *J. Clin. Endocrinol. Metab.* **81**, 745-51.
- Grimm, C., Kohli, J., Murray, J. and Maundrell, K. (1988) Genetic engineering of *Schizosaccharomyces pombe*: a system for gene disruption and replacement using the *ura4* gene as a selectable marker. *Mol. Gen. Genet.* **215**, 81-86.

- Hamm, H.E., Deretic, D., Arendt, A., Hargrave, P.A., Koenig, B. and Hofmann, K.P. (1988) Site of G protein binding to rhodopsin mapped with synthetic peptides from the α -subunit. *Science* **241**, 832-835.
- Hildebrandt, V., Fendler, K., Heberle, J., Hofmann, A., Bamberg, E. and Büldt, G. (1993) Bacteriorhodopsin expressed in *Schizosaccharomyces pombe* pumps protons through the plasma membrane. *Proc. Natl. Acad. Sci. USA* **90**, 3578-3582.
- Hirsch, J.P., Dietzel, C. and Kurjan, J. (1991) The carboxyl terminus of Scg1, the G α subunit involved in yeast mating, is implicated in interactions with the pheromone receptors. *Genes Dev.* **5**, 467-474.
- Jones, R.H., Moreno, S., Nurse, P. and Jones, N.J. (1988) Expression of the SV40 promoter in fission yeast: Identification and characterisation of an AP-1-like factor. *Cell* **53**, 659-667.
- Kajkowski, E.M., Price, L.A., Pausch, M.H., Young, K.H. and Ozenberger, B.A. (1997) Investigation of growth hormone releasing hormone receptor structure and activity using yeast expression technologies. *J. Receptor Signal Transduction Res.* **17**, 293-303.
- King, K., Dohlman, H.G., Thorner, J., Caron, M.G. and Lefkowitz, R.J. (1990) Control of yeast mating signal transduction by a mammalian β 2-adrenergic receptor and Gs α -subunit. *Science* **250**, 121-123.
- Klar, A.J.S. and Bonaduce, M.J. (1991) *swi6*, a gene required for mating-type switching, prohibits meiotic recombination in the *mat2-mat3* "cold-spot" of fission yeast. *Genetics* **129**, 1033-1042.
- Klar, A.J.S. and Miglio, L.M. (1986) Initiation of meiotic recombination by double-strand DNA breaks in *Schizosaccharomyces pombe*. *Cell* **46**, 725-731.
- Klein, C., Paul, J.I., Sauve, K., Schmidt, M.M., Arcangeli, L., Ransom, J., Trueheart, J., Manfredi, J.P., Broach, J.R. and Murphy, A.J. (1998) Identification of surrogate agonists for the human FPRL-1 receptor by autocrine selection in yeast. *Nature Biotech.* **16**, 1334-1337.
- Komatsuzaki, K., Murayama, Y., Giambarella, U., Ogata, E., Seino, S. and Nishimoto, I. (1997) A novel system that reports the G-proteins linked to a given receptor: a study of the type 3 somatostatin receptor. *FEBS Letts* **406**, 165-170.

- Ladds, G., Rasmussen, E.M., Young, T., Nielsen, O. and Davey, J. (1996) The *sxa2*-dependent inactivation of the P-factor mating pheromone in the fission yeast *Schizosaccharomyces pombe*. *Mol. Microbiol.* **20**, 35-42.
- Liu, J., Conklin, B.R., Blin, N., Yun, J. and Wess, J. (1995) Identification of a receptor/G-protein contact site critical for signaling specificity and G-protein activation. *Proc. Natl. Acad. Sci. USA* **92**, 11642-11646.
- Maeda, T., Mochizuki, N. and Yamamoto, M. (1990) Adenylyl cyclase is dispensable for vegetative cell growth in the fission yeast *Schizosaccharomyces pombe*. *Proc. Natl. Acad. Sci. USA* **87**, 7814-7818.
- Milligan, G., Marshal, F. and Rees, S. (1996) $G\alpha_{16}$ as a universal G protein adapter: implications for agonist screening strategies. *Trends Pharmacol. Sci.* **17**, 235-237.
- Murakami, S., Bodley, J.W. and Livingston, D.M. (1982) *Saccharomyces cerevisiae* spheroplasts are sensitive to the action of diphtheria toxin. *Mol. Cell. Biol.* **2**, 588-592.
- Olesnicky, N.S., Brown, A.J., Dowell, S.J. and Casselton, L.A. (1999) A constitutively active G-protein-coupled receptor causes mating self-incompatibility in the mushroom *Coprinus*. *EMBO J.* **18**, 2756-2763.
- Osumi, M. (1998) The ultrastructure of yeast: cell wall structure and formation. *Micron* **29**, 207-233.
- Palm, D., Munch, G., Malek, D., Dees, C. and Heckman, M. (1990) Identification of a Gs protein coupling domain to the beta-adrenoreceptor using site-specific synthetic peptides. Carboxyl terminus of $G\alpha_s$ is involved in coupling to the β -adrenoreceptors. *FEBS Letts* **261**, 294-298.
- Price, L.A., Kajkowski, E.M., Hadcock, J.R., Ozenberger, B.A. and Pausch, M.H. (1995) Functional coupling of a mammalian somatostatin receptor to the yeast pheromone response pathway. *Mol. Cell. Biol.* **15**, 6188-6195.
- Price, L.A., Strnad, J., Pausch, M.H. and Hadcock, J.R. (1996) Pharmacological characterisation of the rat A_{2A} adenosine receptor functionally coupled to the yeast pheromone response pathway. *Mol. Pharmacol.* **50**, 829-837.
- Rasenick, M.M., Watanabe, M., Lazarevic, M.B., Hatta, S. and Hamm, H.E. (1994) Synthetic peptides as probes for G protein function. *J. Biol. Chem.* **269**, 21519-21525.
- Reiländer, H. and Weib, H.M. (1998) Production of G-protein-coupled receptors in yeast. *Current Opinion Biotechnol.* **9**, 510-517.

- Sander, P., Grünewald, S., Reiländer, H. and Michel, H. (1994) Expression of the human D_{2S} dopamine receptor in the yeasts *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe* - a comparative study. *FEBS Letts.* **344**, 41-46.
- Smits, G.J., Kapteyn, J.C., van den Ende, H. and Klis, F.M. (1999) Cell wall dynamics in yeast. *Current Opinion Microbiol.* **2**, 348-352.
- Swift, S., Sheridan, P.J., Covic, L. and Kulipulos, A. (2000) PAR1 thrombin receptor-G protein interactions. Separation of binding and coupling determinants in the G α subunit. *J. Biol. Chem.* **275**, 2627-2635.
- Vale W, Spiess J, Rivier C, Rivier J. (1981) Characterization of a 41-residue ovine hypothalamic peptide that stimulates secretion of corticotropin and β -endorphin. *Science* **213**, 1394-1397.
- Voyno-Yasenetskaya, T., Conklin, B.R., Gilbert, R.L., Hooley, R., Bourne, H.R. and Barber, D.L. (1994) G α 13 stimulates Na-H exchange. *J. Biol. Chem.* **269**, 4721-4724.
- Watson, P., Davis, K., Didmon, M., Broad, P. and Davey, J. (1999) An RGS protein regulates the pheromone response in the fission yeast *Schizosaccharomyces pombe*. *Mol. Microbiol.* **33**, 623-634.

Claims

1. A *Schizosaccharomyces pombe* (*Sz. pombe*) yeast cell comprising:
 - (i) a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway which is derepressed during the mitotic phase of cell growth;
 - (ii) a reporter system for reporting a signal mediated by the GPCR-regulated signalling pathway; and wherein:
 - (a) the GPCR is heterologous, and/or
 - (b) the reporter system comprises a reporter gene and a promoter, the reporter gene being operatively linked to the promoter, and the promoter being regulatable by the GPCR, at least one of the reporter gene and the promoter being heterologous.
2. A yeast cell as claimed in claim 1, wherein the GPCR-regulated signalling pathway is derepressed during the mitotic phase of cell growth by disruption of a nutritional control pathway.
3. A yeast cell according to claim 1 or claim 2, wherein the yeast cell is adenylate cyclase deficient.
4. A yeast cell according to any preceding claim which comprises a mutated *pat1* gene, or in which the endogenous *pat1* gene has been deleted.
5. A yeast cell as claimed in any one of claims 1 to 4, wherein the reporter system comprises a reporter gene and a promoter, the reporter gene being operatively linked to the promoter, and the promoter being regulatable by the GPCR, at least one of the reporter gene and the promoter being heterologous.
6. A yeast cell as claimed in claim 5, wherein the reporter gene is heterologous.
7. A yeast cell according to any preceding claim, wherein the GPCR is a mammalian GPCR.

8. A yeast cell according to any preceding claim comprising a heterologous or chimeric G-protein subunit.
9. A yeast cell according to claim 8, wherein the chimeric G-protein subunit is a $G\alpha$ -transplant.
10. A yeast cell according to claim 9 wherein the $G\alpha$ -transplant is selected from the following transplants:
 - $G\alpha_q$ (SEQ ID 17)
 - $G\alpha_s$ (SEQ ID 16)
 - $G\alpha_o$ (SEQ ID 18)
 - $G\alpha_i2$ (SEQ ID 19)
 - $G\alpha_i3$ (SEQ ID 20)
 - $G\alpha_z$ (SEQ ID 21)
 - $G\alpha_{12}$ (SEQ ID 22)
 - $G\alpha_{13}$ (SEQ ID 23)
 - $G\alpha_{14}$ (SEQ ID 24) and
 - $G\alpha_{16}$ (SEQ ID 25)
11. A yeast cell according to claim 5, wherein the reporter system is regulated by yeast mating pheromone binding to its GPCR.
12. A yeast cell according to claim 11, wherein the yeast mating pheromone is P-factor pheromone.
13. A yeast cell according to claim 12, wherein the reporter system is operatively linked to an *sxa2* promoter or a homologue or analogue thereof.
14. A yeast cell according to any preceding claim wherein the reporter system is integrated into the chromosome of the yeast cell.

15. A yeast cell according to any preceding claim, wherein the yeast cell has a stable mating type.
16. A yeast cell according to any preceding claim wherein the yeast cell is *rgs1* deficient.
17. A yeast cell according to any preceding claim, wherein the yeast cell is *pmp1* deficient.
18. A yeast cell according to any preceding claim, wherein the yeast cell is *sxa2* deficient.
19. A yeast cell according to claim 18, wherein at least a part of the endogenous *sxa2* gene has been deleted.
20. A yeast cell according to claim 19, wherein the reporter gene replaces the deleted *sxa2* gene.
21. A yeast cell according to any preceding claim wherein the reporter gene encodes orotidine-5'-phosphate decarboxylase (the product of the *Sz. pombe ura4* gene), β -galactosidase (the product of the bacterial *lacZ* gene), a β -lactamase, aequorin, green fluorescent protein or luciferase.
22. A yeast cell which is *Schizosaccharomyces pombe* strain JY546 deposited as accession number NCYC2984.
23. A yeast cell according to any preceding claim additionally comprising one or more compounds, to be assayed for their effect on GPCR-regulated expression of the reporter system, or a DNA molecule encoding one or more peptides or proteins to be assayed.

24. A yeast cell according to claim 23, comprising one or more plasmids encoding the or each peptide or protein.
25. A yeast cell according to claim 24, wherein the DNA encoding peptide or protein is transcribed under the control of a thiamine-regulated *nmt1* promoter.
26. A yeast cell according to claim 23 or claim 24, wherein the peptide or protein is of a random sequence.
27. A yeast cell according to any preceding claim, wherein the yeast cell expresses an orphan receptor as the GPCR and the reporter system is regulatable by the orphan receptor.
28. An isolated nucleic acid molecule comprising an *sxa2* promoter, or a homologue or analogue thereof, operatively linked to an exogenous reporter gene.
29. An isolated nucleic acid molecule according to claim 28, wherein the reporter system encodes orotidine-5'-phosphate decarboxylase (the product of the *Sz. pombe ura4* gene), β -galactosidase (the product of the bacterial *lacZ* gene), a β -lactamase, aequorin, green fluorescent protein or luciferase.
30. An isolated nucleic acid molecule encoding a G α -transplant having a nucleic acid sequence selected from:
- G α q (SEQ ID 17)
 - G α s (SEQ ID 16)
 - G α o (SEQ ID 18)
 - G α i2 (SEQ ID 19)
 - G α i3 (SEQ ID 20)
 - G α z (SEQ ID 21)
 - G α 12 (SEQ ID 22)
 - G α 13 (SEQ ID 23)
 - G α 14 (SEQ ID 24) and

Gα16 (SEQ ID 25);

or which differs from the one or more of the sequences due to degeneracy in the genetic code.

31. Use of:

(a) a *Schizosaccharomyces pombe* (*Sz. pombe*) yeast cell comprising:

(i) a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway which is derepressed during the mitotic phase of cell growth;

(ii) a reporter system for reporting a signal mediated by the GPCR-regulated signalling pathway; or

(b) an isolated nucleic acid molecule according to any one of claims 28 to 30 to study GPCR-regulated activity.

32. An assay comprising the use of a yeast cell or isolated DNA molecule according to any one of claims 1 to 31.

33. A method of determining the effect of a compound, on GPCR-regulated activity comprising the steps of:

(i) providing a *Schizosaccharomyces pombe* (*Sz. pombe*) yeast cell comprising:

(a) a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway which is derepressed during the mitotic phase of cell growth;

(b) a reporter system for reporting a signal mediated by the GPCR-regulated signalling pathway;

(ii) introducing the compound, to the yeast cell; and

(iii) noting the output of the reporter system.

34. Use of a yeast cell according to claim 27 to identify compounds, which affect the ability of the orphan GPCR to regulate the reporter system.

35. Use of a *Schizosaccharomyces pombe* (*Sz. pombe*) yeast cell comprising:

(i) a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway which is derepressed during the mitotic phase of cell growth;

(ii) a reporter system for reporting a signal mediated by the GPCR-regulated signalling pathway;
to identify a regulator or a mutant of a component of a GPCR-regulated pathway.

36. A method of identifying a reagent that modulates GPCR-regulated signalling pathways comprising:

(i) providing a *Schizosaccharomyces pombe* (*Sz. pombe*) yeast cell comprising:

(a) a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway which is derepressed during the mitotic phase of cell growth;

(b) a reporter system for reporting a signal mediated by the GPCR-regulated signalling pathway;

(ii) producing a random peptide within the yeast cell; and

(iii) measuring an amount of reporter activity produced.

37. A compound capable of modulating GPCR activity identified by a method according to claim 33 or claim 36.

38. A method of determining whether a G-Protein Coupled Receptor (GPCR) is coupled to a cell signalling pathway, the method comprising comparing the ligand-independent reporter system output of a *Schizosaccharomyces pombe* (*Sz. pombe*) yeast cell comprising:

(i) a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway which is derepressed during the mitotic phase of cell growth;

(ii) a reporter system for reporting a signal mediated by the GPCR-regulated signalling pathway;

with the reporter system output of a reference cell which lacks a functional GPCR.

39. A *Schizosaccharomyces pombe* (*Sz. pombe*) yeast cell comprising:

(i) a G-Protein Coupled Receptor (GPCR)-regulated signalling pathway which is derepressed during the mitotic phase of cell growth, wherein the GPCR is absent or otherwise rendered non-functional;

45

(ii) a reporter system for reporting a signal mediated by the GPCR-regulated signalling pathway.

40. An assay kit comprising a yeast cell or isolated nucleic acid molecule as defined in any one of claims 1 to 30 and 39.

Figure 1




Figure 2




Figure 3

3/16




Figure 4

•

MKITAVIALLESLAAASPIPVADPGVVSVSYSYADFLRVYQSWNTFANPDRPNLKKREFEAAPAKTYA
DFLAYQSWNTFVNPDNRPNLKKREFEAAPEKSYADFLRAYHSWNTFVNPDNRPNLKKREFEAAPAKTYA
DFLAYQSWNTFVNPDNRPNLKKRTEDEENEEDEEYYRFLQFYIMTVPENSTITDVNITAKFES

Figure 5

5/16




Figure 6

MKITAVIALLFSLAASPVPVADPGVVSVSKSYADEFLRVYQSWNTFANPDRPNLIKRT
EEDEENEEDEEYRFLQFYIMTVPENSTITDVNITAKFES





Figure 7B

MKITAVIALLFSLAAASPIPVADPGVSVSKnnnnnnnnnnnnnnnnnnnnnnnnKRR
TEEDEENEEDEEYRFLQFYIMTVPENSTITDVTAKFES

Figure 8




Figure 9




Figure 10






Figure 11



Sxa2>Ura4 reporter strain.
MMM - Uracil
Random mutations introduced into rgs1

13/16

Figure 12



Repressed Induced

*Inducible expression of P-factor
Uses nmt1 promoter to express monoP construct
Sxa2> Ura4 reporter strain
DMM - Uracil*

Figure 13




Figure 14





Figure 15



SEQUENCE LISTING

<110> Septegen Ltd

<120> Yeast-Based Assay

<130> DE/p701948PCT

<140>

<141>

<150> GB 0030038.4

<151> 2000-12-08

<160> 53

<170> PatentIn Ver. 2.1

<210> 1

<211> 407

<212> PRT

<213> Schizosaccharomyces pombe

<400> 1

Met Gly Cys Met Ser Ser Lys Tyr Ala Asp Thr Ser Gly Gly Glu Val
1 5 10 15

Ile Gln Lys Lys Leu Ser Asp Thr Gln Thr Ser Asn Ser Ser Thr Thr
20 25 30

Gly Ser Gln Asn Ala Arg Val Pro Val Leu Glu Asn Trp Leu Asn Ile
35 40 45

Val Leu Arg Gly Lys Pro Gln Asn Val Glu Ser Ser Gly Val Arg Val
50 55 60

Lys Gly Asn Ser Thr Ser Gly Gly Asn Asp Ile Lys Val Leu Leu Leu
65 70 75 80

Gly Ala Gly Asp Ser Gly Lys Thr Thr Ile Met Lys Gln Met Arg Leu
85 90 95

Leu Tyr Ser Pro Gly Phe Ser Gln Val Val Arg Lys Gln Tyr Arg Val
100 105 110

Met Ile Phe Glu Asn Ile Ile Ser Ser Leu Cys Leu Leu Leu Glu Ala
115 120 125

Met Asp Asn Ser Asn Val Ser Leu Leu Pro Glu Asn Glu Lys Tyr Arg
 130 135 140

Ala Val Ile Leu Arg Lys His Thr Ser Gln Pro Asn Glu Pro Phe Ser
 145 150 155 160

Pro Glu Ile Tyr Glu Ala Val His Ala Leu Thr Leu Asp Thr Lys Leu
 165 170 175

Arg Thr Val Gln Ser Cys Gly Thr Asn Leu Ser Leu Leu Asp Asn Phe
 180 185 190

Tyr Tyr Tyr Gln Asp His Ile Asp Arg Ile Phe Asp Pro Gln Tyr Ile
 195 200 205

Pro Ser Asp Gln Asp Ile Leu His Cys Arg Ile Lys Thr Thr Gly Ile
 210 215 220

Ser Glu Glu Thr Phe Leu Leu Asn Arg His His Tyr Arg Phe Phe Asp
 225 230 235 240

Val Gly Gly Gln Arg Ser Glu Arg Arg Lys Trp Ile His Cys Phe Glu
 245 250 255

Asn Val Thr Ala Leu Leu Phe Leu Val Ser Leu Ala Gly Tyr Asp Gln
 260 265 270

Cys Leu Val Glu Asp Asn Ser Gly Asn Gln Met Gln Glu Ala Leu Leu
 275 280 285

Leu Trp Asp Ser Ile Cys Asn Ser Ser Trp Phe Ser Glu Ser Ala Met
 290 295 300

Ile Leu Phe Leu Asn Lys Leu Asp Leu Phe Lys Arg Lys Val His Ile
 305 310 315 320

Ser Pro Ile Gln Lys His Phe Pro Asp Tyr Gln Glu Val Gly Ser Thr
 325 330 335

Pro Thr Phe Val Gln Thr Gln Cys Pro Leu Ala Asp Asn Ala Val Arg
 340 345 350

Ser Gly Met Tyr Tyr Phe Tyr Leu Lys Phe Glu Ser Leu Asn Arg Ile
 355 360 365

Ala Ser Arg Ser Cys Tyr Cys His Phe Thr Thr Ala Thr Asp Thr Ser
 370 375 380

Leu Leu Gln Arg Val Met Val Ser Val Gln Asp Thr Ile Met Ser Asn
 385 390 395 400

Asn Leu Gln Ser Leu Met Phe
 405

<210> 2

<211> 407

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: G alpha
 transplant

<400> 2

Met Gly Cys Met Ser Ser Lys Tyr Ala Asp Thr Ser Gly Gly Glu Val
 1 5 10 15

Ile Gln Lys Lys Leu Ser Asp Thr Gln Thr Ser Asn Ser Ser Thr Thr
 20 25 30

Gly Ser Gln Asn Ala Arg Val Pro Val Leu Glu Asn Trp Leu Asn Ile
 35 40 45

Val Leu Arg Gly Lys Pro Gln Asn Val Glu Ser Ser Gly Val Arg Val
 50 55 60

Lys Gly Asn Ser Thr Ser Gly Gly Asn Asp Ile Lys Val Leu Leu Leu
 65 70 75 80

Gly Ala Gly Asp Ser Gly Lys Thr Thr Ile Met Lys Gln Met Arg Leu
 85 90 95

Leu Tyr Ser Pro Gly Phe Ser Gln Val Val Arg Lys Gln Tyr Arg Val
 100 105 110

Met Ile Phe Glu Asn Ile Ile Ser Ser Leu Cys Leu Leu Leu Glu Ala
 115 120 125

Met Asp Asn Ser Asn Val Ser Leu Leu Pro Glu Asn Glu Lys Tyr Arg
 130 135 140

Ala Val Ile Leu Arg Lys His Thr Ser Gln Pro Asn Glu Pro Phe Ser
 145 150 155 160

Pro Glu Ile Tyr Glu Ala Val His Ala Leu Thr Leu Asp Thr Lys Leu
 165 170 175

Arg Thr Val Gln Ser Cys Gly Thr Asn Leu Ser Leu Leu Asp Asn Phe
 180 185 190

Tyr Tyr Tyr Gln Asp His Ile Asp Arg Ile Phe Asp Pro Gln Tyr Ile
 195 200 205

Pro Ser Asp Gln Asp Ile Leu His Cys Arg Ile Lys Thr Thr Gly Ile
 210 215 220

Ser Glu Glu Thr Phe Leu Leu Asn Arg His His Tyr Arg Phe Phe Asp
 225 230 235 240

Val Gly Gly Gln Arg Ser Glu Arg Arg Lys Trp Ile His Cys Phe Glu
 245 250 255

Asn Val Thr Ala Leu Leu Phe Leu Val Ser Leu Ala Gly Tyr Asp Gln
 260 265 270

Cys Leu Val Glu Asp Asn Ser Gly Asn Gln Met Gln Glu Ala Leu Leu
 275 280 285

Leu Trp Asp Ser Ile Cys Asn Ser Ser Trp Phe Ser Glu Ser Ala Met
 290 295 300

Ile Leu Phe Leu Asn Lys Leu Asp Leu Phe Lys Arg Lys Val His Ile
 305 310 315 320

Ser Pro Ile Gln Lys His Phe Pro Asp Tyr Gln Glu Val Gly Ser Thr
 325 330 335

Pro Thr Phe Val Gln Thr Gln Cys Pro Leu Ala Asp Asn Ala Val Arg
 340 345 350

Ser Gly Met Tyr Tyr Phe Tyr Leu Lys Phe Glu Ser Leu Asn Arg Ile
 355 360 365

Ala Ser Arg Ser Cys Tyr Cys His Phe Thr Thr Ala Thr Asp Thr Ser
 370 375 380

Leu Leu Gln Arg Val Met Val Ser Val Gln Asp Thr Ile Met Ser Asn
 385 390 395 400

Asn Leu Gln Tyr Glu Leu Leu
 405

<210> 3

<211> 407

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: G alpha
transplant

<400> 3

Met Gly Cys Met Ser Ser Lys Tyr Ala Asp Thr Ser Gly Gly Glu Val
1 5 10 15

Ile Gln Lys Lys Leu Ser Asp Thr Gln Thr Ser Asn Ser Ser Thr Thr
20 25 30

Gly Ser Gln Asn Ala Arg Val Pro Val Leu Glu Asn Trp Leu Asn Ile
35 40 45

Val Leu Arg Gly Lys Pro Gln Asn Val Glu Ser Ser Gly Val Arg Val
50 55 60

Lys Gly Asn Ser Thr Ser Gly Gly Asn Asp Ile Lys Val Leu Leu Leu
65 70 75 80

Gly Ala Gly Asp Ser Gly Lys Thr Thr Ile Met Lys Gln Met Arg Leu
85 90 95

Leu Tyr Ser Pro Gly Phe Ser Gln Val Val Arg Lys Gln Tyr Arg Val
100 105 110

Met Ile Phe Glu Asn Ile Ile Ser Ser Leu Cys Leu Leu Leu Glu Ala
115 120 125

Met Asp Asn Ser Asn Val Ser Leu Leu Pro Glu Asn Glu Lys Tyr Arg
130 135 140

Ala Val Ile Leu Arg Lys His Thr Ser Gln Pro Asn Glu Pro Phe Ser
145 150 155 160

Pro Glu Ile Tyr Glu Ala Val His Ala Leu Thr Leu Asp Thr Lys Leu
165 170 175

Arg Thr Val Gln Ser Cys Gly Thr Asn Leu Ser Leu Leu Asp Asn Phe
180 185 190

Tyr Tyr Tyr Gln Asp His Ile Asp Arg Ile Phe Asp Pro Gln Tyr Ile
 195 200 205

Pro Ser Asp Gln Asp Ile Leu His Cys Arg Ile Lys Thr Thr Gly Ile
 210 215 220

Ser Glu Glu Thr Phe Leu Leu Asn Arg His His Tyr Arg Phe Phe Asp
 225 230 235 240

Val Gly Gly Gln Arg Ser Glu Arg Arg Lys Trp Ile His Cys Phe Glu
 245 250 255

Asn Val Thr Ala Leu Leu Phe Leu Val Ser Leu Ala Gly Tyr Asp Gln
 260 265 270

Cys Leu Val Glu Asp Asn Ser Gly Asn Gln Met Gln Glu Ala Leu Leu
 275 280 285

Leu Trp Asp Ser Ile Cys Asn Ser Ser Trp Phe Ser Glu Ser Ala Met
 290 295 300

Ile Leu Phe Leu Asn Lys Leu Asp Leu Phe Lys Arg Lys Val His Ile
 305 310 315 320

Ser Pro Ile Gln Lys His Phe Pro Asp Tyr Gln Glu Val Gly Ser Thr
 325 330 335

Pro Thr Phe Val Gln Thr Gln Cys Pro Leu Ala Asp Asn Ala Val Arg
 340 345 350

Ser Gly Met Tyr Tyr Phe Tyr Leu Lys Phe Glu Ser Leu Asn Arg Ile
 355 360 365

Ala Ser Arg Ser Cys Tyr Cys His Phe Thr Thr Ala Thr Asp Thr Ser
 370 375 380

Leu Leu Gln Arg Val Met Val Ser Val Gln Asp Thr Ile Met Ser Asn
 385 390 395 400

Asn Leu Glu Tyr Asn Leu Val
 405

<210> 4

<211> 407

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: G alpha
transplant

<400> 4

Met Gly Cys Met Ser Ser Lys Tyr Ala Asp Thr Ser Gly Gly Glu Val
1 5 10 15

Ile Gln Lys Lys Leu Ser Asp Thr Gln Thr Ser Asn Ser Ser Thr Thr
20 25 30

Gly Ser Gln Asn Ala Arg Val Pro Val Leu Glu Asn Trp Leu Asn Ile
35 40 45

Val Leu Arg Gly Lys Pro Gln Asn Val Glu Ser Ser Gly Val Arg Val
50 55 60

Lys Gly Asn Ser Thr Ser Gly Gly Asn Asp Ile Lys Val Leu Leu Leu
65 70 75 80

Gly Ala Gly Asp Ser Gly Lys Thr Thr Ile Met Lys Gln Met Arg Leu
85 90 95

Leu Tyr Ser Pro Gly Phe Ser Gln Val Val Arg Lys Gln Tyr Arg Val
100 105 110

Met Ile Phe Glu Asn Ile Ile Ser Ser Leu Cys Leu Leu Glu Ala
115 120 125

Met Asp Asn Ser Asn Val Ser Leu Leu Pro Glu Asn Glu Lys Tyr Arg
130 135 140

Ala Val Ile Leu Arg Lys His Thr Ser Gln Pro Asn Glu Pro Phe Ser
145 150 155 160

Pro Glu Ile Tyr Glu Ala Val His Ala Leu Thr Leu Asp Thr Lys Leu
165 170 175

Arg Thr Val Gln Ser Cys Gly Thr Asn Leu Ser Leu Leu Asp Asn Phe
180 185 190

Tyr Tyr Tyr Gln Asp His Ile Asp Arg Ile Phe Asp Pro Gln Tyr Ile
195 200 205

Pro Ser Asp Gln Asp Ile Leu His Cys Arg Ile Lys Thr Thr Gly Ile
210 215 220

Ser Glu Glu Thr Phe Leu Leu Asn Arg His His Tyr Arg Phe Phe Asp
225 230 235 240

Val Gly Gly Gln Arg Ser Glu Arg Arg Lys Trp Ile His Cys Phe Glu
245 250 255

Asn Val Thr Ala Leu Leu Phe Leu Val Ser Leu Ala Gly Tyr Asp Gln
260 265 270

Cys Leu Val Glu Asp Asn Ser Gly Asn Gln Met Gln Glu Ala Leu Leu
275 280 285

Leu Trp Asp Ser Ile Cys Asn Ser Ser Trp Phe Ser Glu Ser Ala Met
290 295 300

Ile Leu Phe Leu Asn Lys Leu Asp Leu Phe Lys Arg Lys Val His Ile
305 310 315 320

Ser Pro Ile Gln Lys His Phe Pro Asp Tyr Gln Glu Val Gly Ser Thr
325 330 335

Pro Thr Phe Val Gln Thr Gln Cys Pro Leu Ala Asp Asn Ala Val Arg
340 345 350

Ser Gly Met Tyr Tyr Phe Tyr Leu Lys Phe Glu Ser Leu Asn Arg Ile
355 360 365

Ala Ser Arg Ser Cys Tyr Cys His Phe Thr Thr Ala Thr Asp Thr Ser
370 375 380

Leu Leu Gln Arg Val Met Val Ser Val Gln Asp Thr Ile Met Ser Asn
385 390 395 400

Asn Leu Gly Cys Gly Leu Tyr
405

<210> 5

<211> 407

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: G alpha
transplant

<400> 5

Met Gly Cys Met Ser Ser Lys Tyr Ala Asp Thr Ser Gly Gly Glu Val
 1 5 10 15
 Ile Gln Lys Lys Leu Ser Asp Thr Gln Thr Ser Asn Ser Ser Thr Thr
 20 25 30
 Gly Ser Gln Asn Ala Arg Val Pro Val Leu Glu Asn Trp Leu Asn Ile
 35 40 45
 Val Leu Arg Gly Lys Pro Gln Asn Val Glu Ser Ser Gly Val Arg Val
 50 55 60
 Lys Gly Asn Ser Thr Ser Gly Gly Asn Asp Ile Lys Val Leu Leu Leu
 65 70 75 80
 Gly Ala Gly Asp Ser Gly Lys Thr Thr Ile Met Lys Gln Met Arg Leu
 85 90 95
 Leu Tyr Ser Pro Gly Phe Ser Gln Val Val Arg Lys Gln Tyr Arg Val
 100 105 110
 Met Ile Phe Glu Asn Ile Ile Ser Ser Leu Cys Leu Leu Leu Glu Ala
 115 120 125
 Met Asp Asn Ser Asn Val Ser Leu Leu Pro Glu Asn Glu Lys Tyr Arg
 130 135 140
 Ala Val Ile Leu Arg Lys His Thr Ser Gln Pro Asn Glu Pro Phe Ser
 145 150 155 160
 Pro Glu Ile Tyr Glu Ala Val His Ala Leu Thr Leu Asp Thr Lys Leu
 165 170 175
 Arg Thr Val Gln Ser Cys Gly Thr Asn Leu Ser Leu Leu Asp Asn Phe
 180 185 190
 Tyr Tyr Tyr Gln Asp His Ile Asp Arg Ile Phe Asp Pro Gln Tyr Ile
 195 200 205
 Pro Ser Asp Gln Asp Ile Leu His Cys Arg Ile Lys Thr Thr Gly Ile
 210 215 220
 Ser Glu Glu Thr Phe Leu Leu Asn Arg His His Tyr Arg Phe Phe Asp
 225 230 235 240
 Val Gly Gly Gln Arg Ser Glu Arg Arg Lys Trp Ile His Cys Phe Glu
 245 250 255

Asn Val Thr Ala Leu Leu Phe Leu Val Ser Leu Ala Gly Tyr Asp Gln
 260 265 270

Cys Leu Val Glu Asp Asn Ser Gly Asn Gln Met Gln Glu Ala Leu Leu
 275 280 285

Leu Trp Asp Ser Ile Cys Asn Ser Ser Trp Phe Ser Glu Ser Ala Met
 290 295 300

Ile Leu Phe Leu Asn Lys Leu Asp Leu Phe Lys Arg Lys Val His Ile
 305 310 315 320

Ser Pro Ile Gln Lys His Phe Pro Asp Tyr Gln Glu Val Gly Ser Thr
 325 330 335

Pro Thr Phe Val Gln Thr Gln Cys Pro Leu Ala Asp Asn Ala Val Arg
 340 345 350

Ser Gly Met Tyr Tyr Phe Tyr Leu Lys Phe Glu Ser Leu Asn Arg Ile
 355 360 365

Ala Ser Arg Ser Cys Tyr Cys His Phe Thr Thr Ala Thr Asp Thr Ser
 370 375 380

Leu Leu Gln Arg Val Met Val Ser Val Gln Asp Thr Ile Met Ser Asn
 385 390 395 400

Asn Leu Asp Cys Gly Leu Phe
 405

<210> 6

<211> 407

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: G alpha
 transplant

<400> 6

Met Gly Cys Met Ser Ser Lys Tyr Ala Asp Thr Ser Gly Gly Glu Val
 1 5 10 15

Ile Gln Lys Lys Leu Ser Asp Thr Gln Thr Ser Asn Ser Ser Thr Thr
 20 25 30

Gly Ser Gln Asn Ala Arg Val Pro Val Leu Glu Asn Trp Leu Asn Ile
 35 40 45

Val Leu Arg Gly Lys Pro Gln Asn Val Glu Ser Ser Gly Val Arg Val
 50 55 60

Lys Gly Asn Ser Thr Ser Gly Gly Asn Asp Ile Lys Val Leu Leu Leu
 65 70 75 80

Gly Ala Gly Asp Ser Gly Lys Thr Thr Ile Met Lys Gln Met Arg Leu
 85 90 95

Leu Tyr Ser Pro Gly Phe Ser Gln Val Val Arg Lys Gln Tyr Arg Val
 100 105 110

Met Ile Phe Glu Asn Ile Ile Ser Ser Leu Cys Leu Leu Leu Glu Ala
 115 120 125

Met Asp Asn Ser Asn Val Ser Leu Leu Pro Glu Asn Glu Lys Tyr Arg
 130 135 140

Ala Val Ile Leu Arg Lys His Thr Ser Gln Pro Asn Glu Pro Phe Ser
 145 150 155 160

Pro Glu Ile Tyr Glu Ala Val His Ala Leu Thr Leu Asp Thr Lys Leu
 165 170 175

Arg Thr Val Gln Ser Cys Gly Thr Asn Leu Ser Leu Leu Asp Asn Phe
 180 185 190

Tyr Tyr Tyr Gln Asp His Ile Asp Arg Ile Phe Asp Pro Gln Tyr Ile
 195 200 205

Pro Ser Asp Gln Asp Ile Leu His Cys Arg Ile Lys Thr Thr Gly Ile
 210 215 220

Ser Glu Glu Thr Phe Leu Leu Asn Arg His His Tyr Arg Phe Phe Asp
 225 230 235 240

Val Gly Gly Gln Arg Ser Glu Arg Arg Lys Trp Ile His Cys Phe Glu
 245 250 255

Asn Val Thr Ala Leu Leu Phe Leu Val Ser Leu Ala Gly Tyr Asp Gln
 260 265 270

Cys Leu Val Glu Asp Asn Ser Gly Asn Gln Met Gln Glu Ala Leu Leu
 275 280 285

12/55

Leu Trp Asp Ser Ile Cys Asn Ser Ser Trp Phe Ser Glu Ser Ala Met
 290 295 300

Ile Leu Phe Leu Asn Lys Leu Asp Leu Phe Lys Arg Lys Val His Ile
 305 310 315 320

Ser Pro Ile Gln Lys His Phe Pro Asp Tyr Gln Glu Val Gly Ser Thr
 325 330 335

Pro Thr Phe Val Gln Thr Gln Cys Pro Leu Ala Asp Asn Ala Val Arg
 340 345 350

Ser Gly Met Tyr Tyr Phe Tyr Leu Lys Phe Glu Ser Leu Asn Arg Ile
 355 360 365

Ala Ser Arg Ser Cys Tyr Cys His Phe Thr Thr Ala Thr Asp Thr Ser
 370 375 380

Leu Leu Gln Arg Val Met Val Ser Val Gln Asp Thr Ile Met Ser Asn
 385 390 395 400

Asn Leu Glu Cys Gly Leu Tyr
 405

<210> 7

<211> 407

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: G alpha
 transplant

<400> 7

Met Gly Cys Met Ser Ser Lys Tyr Ala Asp Thr Ser Gly Gly Glu Val
 1 5 10 15

Ile Gln Lys Lys Leu Ser Asp Thr Gln Thr Ser Asn Ser Ser Thr Thr
 20 25 30

Gly Ser Gln Asn Ala Arg Val Pro Val Leu Glu Asn Trp Leu Asn Ile
 35 40 45

Val Leu Arg Gly Lys Pro Gln Asn Val Glu Ser Ser Gly Val Arg Val
 50 55 60

Lys Gly Asn Ser Thr Ser Gly Gly Asn Asp Ile Lys Val Leu Leu Leu
65 70 75 80

Gly Ala Gly Asp Ser Gly Lys Thr Thr Ile Met Lys Gln Met Arg Leu
85 90 95

Leu Tyr Ser Pro Gly Phe Ser Gln Val Val Arg Lys Gln Tyr Arg Val
100 105 110

Met Ile Phe Glu Asn Ile Ile Ser Ser Leu Cys Leu Leu Leu Glu Ala
115 120 125

Met Asp Asn Ser Asn Val Ser Leu Leu Pro Glu Asn Glu Lys Tyr Arg
130 135 140

Ala Val Ile Leu Arg Lys His Thr Ser Gln Pro Asn Glu Pro Phe Ser
145 150 155 160

Pro Glu Ile Tyr Glu Ala Val His Ala Leu Thr Leu Asp Thr Lys Leu
165 170 175

Arg Thr Val Gln Ser Cys Gly Thr Asn Leu Ser Leu Leu Asp Asn Phe
180 185 190

Tyr Tyr Tyr Gln Asp His Ile Asp Arg Ile Phe Asp Pro Gln Tyr Ile
195 200 205

Pro Ser Asp Gln Asp Ile Leu His Cys Arg Ile Lys Thr Thr Gly Ile
210 215 220

Ser Glu Glu Thr Phe Leu Leu Asn Arg His His Tyr Arg Phe Phe Asp
225 230 235 240

Val Gly Gly Gln Arg Ser Glu Arg Arg Lys Trp Ile His Cys Phe Glu
245 250 255

Asn Val Thr Ala Leu Leu Phe Leu Val Ser Leu Ala Gly Tyr Asp Gln
260 265 270

Cys Leu Val Glu Asp Asn Ser Gly Asn Gln Met Gln Glu Ala Leu Leu
275 280 285

Leu Trp Asp Ser Ile Cys Asn Ser Ser Trp Phe Ser Glu Ser Ala Met
290 295 300

Ile Leu Phe Leu Asn Lys Leu Asp Leu Phe Lys Arg Lys Val His Ile
305 310 315 320

Pro Thr Phe Val Gln Thr Gln Cys Pro Leu Ala Asp Asn Ala Val Arg
340 345 350

Ser Gly Met Tyr Tyr Phe Tyr Leu Lys Phe Glu Ser Leu Asn Arg Ile
355 360 365

Ala Ser Arg Ser Cys Tyr Cys His Phe Thr Thr Ala Thr Asp Thr Ser
370 375 380

Leu Leu Gln Arg Val Met Val Ser Val Gln Asp Thr Ile Met Ser Asn
385 390 395 400

Asn Leu Tyr Ile Gly Leu Cys
405

<210> 8

<211> 407

<212> PRT

<213> Artificial Sequence

$\langle 220 \rangle$

<220>
<223> Description of Artificial Sequence: G alpha
transplant

<400> 8

<400> 8
Met Gly Cys Met Ser Ser Lys Tyr Ala Asp Thr Ser Gly Gly Glu Val
1 5 10 15

Ile Gln Lys Lys Leu Ser Asp Thr Gln Thr Ser Asn Ser Ser Thr Thr
20 25 30

Gly Ser Gln Asn Ala Arg Val Pro Val Leu Glu Asn Trp Leu Asn Ile
35 40 45

Val Leu Arg Gly Lys Pro Gln Asn Val Glu Ser Ser Gly Val Arg Val
50 55 60

Lys Gly Asn Ser Thr Ser Gly Gly Asn Asp Ile Lys Val Leu Leu Leu
65 70 75 .80

Gly Ala Gly Asp Ser Gly Lys Thr Thr Ile Met Lys Gln Met Arg Leu
85 90 95

Leu Tyr Ser Pro Gly Phe Ser Gln Val Val Arg Lys Gln Tyr Arg Val
100 105 110

Met Ile Phe Glu Asn Ile Ile Ser Ser Leu Cys Leu Leu Leu Glu Ala
115 120 125

Met Asp Asn Ser Asn Val Ser Leu Leu Pro Glu Asn Glu Lys Tyr Arg
130 135 140

Ala Val Ile Leu Arg Lys His Thr Ser Gln Pro Asn Glu Pro Phe Ser
145 150 155 160

Pro Glu Ile Tyr Glu Ala Val His Ala Leu Thr Leu Asp Thr Lys Leu
165 170 175

Arg Thr Val Gln Ser Cys Gly Thr Asn Leu Ser Leu Leu Asp Asn Phe
180 185 190

Tyr Tyr Tyr Gln Asp His Ile Asp Arg Ile Phe Asp Pro Gln Tyr Ile
195 200 205

Pro Ser Asp Gln Asp Ile Leu His Cys Arg Ile Lys Thr Thr Gly Ile
210 215 220

Ser Glu Glu Thr Phe Leu Leu Asn Arg His His Tyr Arg Phe Phe Asp
225 230 235 240

Val Gly Gly Gln Arg Ser Glu Arg Arg Lys Trp Ile His Cys Phe Glu
245 250 255

Asn Val Thr Ala Leu Leu Phe Leu Val Ser Leu Ala Gly Tyr Asp Gln
260 265 270

Cys Leu Val Glu Asp Asn Ser Gly Asn Gln Met Gln Glu Ala Leu Leu
275 280 285

Leu Trp Asp Ser Ile Cys Asn Ser Ser Trp Phe Ser Glu Ser Ala Met
290 295 300

Ile Leu Phe Leu Asn Lys Leu Asp Leu Phe Lys Arg Lys Val His Ile
305 310 315 320

Ser Pro Ile Gln Lys His Phe Pro Asp Tyr Gln Glu Val Gly Ser Thr
325 330 335

Pro Thr Phe Val Gln Thr Gln Cys Pro Leu Ala Asp Asn Ala Val Arg
340 345 350

Ser Gly Met Tyr Tyr Phe Tyr Leu Lys Phe Glu Ser Leu Asn Arg Ile
 355 360 365

Ala Ser Arg Ser Cys Tyr Cys His Phe Thr Thr Ala Thr Asp Thr Ser
 370 375 380

Leu Leu Gln Arg Val Met Val Ser Val Gln Asp Thr Ile Met Ser Asn
 385 390 395 400

Asn Leu Glu Asn Val Arg Phe
 405

<210> 9

<211> 407

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: G alpha
 transplant

<400> 9

Met Gly Cys Met Ser Ser Lys Tyr Ala Asp Thr Ser Gly Gly Glu Val
 1 5 10 15

Ile Gln Lys Lys Leu Ser Asp Thr Gln Thr Ser Asn Ser Ser Thr Thr
 20 25 30

Gly Ser Gln Asn Ala Arg Val Pro Val Leu Glu Asn Trp Leu Asn Ile
 35 40 45

Val Leu Arg Gly Lys Pro Gln Asn Val Glu Ser Ser Gly Val Arg Val
 50 55 60

Lys Gly Asn Ser Thr Ser Gly Gly Asn Asp Ile Lys Val Leu Leu Leu
 65 70 75 80

Gly Ala Gly Asp Ser Gly Lys Thr Thr Ile Met Lys Gln Met Arg Leu
 85 90 95

Leu Tyr Ser Pro Gly Phe Ser Gln Val Val Arg Lys Gln Tyr Arg Val
 100 105 110

Met Ile Phe Glu Asn Ile Ile Ser Ser Leu Cys Leu Leu Leu Glu Ala
 115 120 125

17/55

Met Asp Asn Ser Asn Val Ser Leu Leu Pro Glu Asn Glu Lys Tyr Arg
130 135 140

Ala Val Ile Leu Arg Lys His Thr Ser Gln Pro Asn Glu Pro Phe Ser
145 150 155 160

Pro Glu Ile Tyr Glu Ala Val His Ala Leu Thr Leu Asp Thr Lys Leu
165 170 175

Arg Thr Val Gln Ser Cys Gly Thr Asn Leu Ser Leu Leu Asp Asn Phe
180 185 190

Tyr Tyr Tyr Gln Asp His Ile Asp Arg Ile Phe Asp Pro Gln Tyr Ile
195 200 205

Pro Ser Asp Gln Asp Ile Leu His Cys Arg Ile Lys Thr Thr Gly Ile
210 215 220

Ser Glu Glu Thr Phe Leu Leu Asn Arg His His Tyr Arg Phe Phe Asp
225 230 235 240

Val Gly Gly Gln Arg Ser Glu Arg Arg Lys Trp Ile His Cys Phe Glu
245 250 255

Asn Val Thr Ala Leu Leu Phe Leu Val Ser Leu Ala Gly Tyr Asp Gln
260 265 270

Cys Leu Val Glu Asp Asn Ser Gly Asn Gln Met Gln Glu Ala Leu Leu
275 280 285

Leu Trp Asp Ser Ile Cys Asn Ser Ser Trp Phe Ser Glu Ser Ala Met
290 295 300

Ile Leu Phe Leu Asn Lys Leu Asp Leu Phe Lys Arg Lys Val His Ile
305 310 315 320

Ser Pro Ile Gln Lys His Phe Pro Asp Tyr Gln Glu Val Gly Ser Thr
325 330 335

Pro Thr Phe Val Gln Thr Gln Cys Pro Leu Ala Asp Asn Ala Val Arg
340 345 350

Ser Gly Met Tyr Tyr Phe Tyr Leu Lys Phe Glu Ser Leu Asn Arg Ile
355 360 365

Ala Ser Arg Ser Cys Tyr Cys His Phe Thr Thr Ala Thr Asp Thr Ser
370 375 380

Leu Leu Gln Arg Val Met Val Ser Val Gln Asp Thr Ile Met Ser Asn
 385 390 395 400

Asn Leu Arg Leu Val Phe Arg
 405

<210> 10

<211> 407

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: G alpha
 transplant

<400> 10

Met Gly Cys Met Ser Ser Lys Tyr Ala Asp Thr Ser Gly Gly Glu Val
 1 5 10 15

Ile Gln Lys Lys Leu Ser Asp Thr Gln Thr Ser Asn Ser Ser Thr Thr
 20 25 30

Gly Ser Gln Asn Ala Arg Val Pro Val Leu Glu Asn Trp Leu Asn Ile
 35 40 45

Val Leu Arg Gly Lys Pro Gln Asn Val Glu Ser Ser Gly Val Arg Val
 50 55 60

Lys Gly Asn Ser Thr Ser Gly Gly Asn Asp Ile Lys Val Leu Leu Leu
 65 70 75 80

Gly Ala Gly Asp Ser Gly Lys Thr Thr Ile Met Lys Gln Met Arg Leu
 85 90 95

Leu Tyr Ser Pro Gly Phe Ser Gln Val Val Arg Lys Gln Tyr Arg Val
 100 105 110

Met Ile Phe Glu Asn Ile Ile Ser Ser Leu Cys Leu Leu Leu Glu Ala
 115 120 125

Met Asp Asn Ser Asn Val Ser Leu Leu Pro Glu Asn Glu Lys Tyr Arg
 130 135 140

Ala Val Ile Leu Arg Lys His Thr Ser Gln Pro Asn Glu Pro Phe Ser
 145 150 155 160

Pro Glu Ile Tyr Glu Ala Val His Ala Leu Thr Leu Asp Thr Lys Leu
 165 170 175

Arg Thr Val Gln Ser Cys Gly Thr Asn Leu Ser Leu Leu Asp Asn Phe
 180 185 190

Tyr Tyr Tyr Gln Asp His Ile Asp Arg Ile Phe Asp Pro Gln Tyr Ile
 195 200 205

Pro Ser Asp Gln Asp Ile Leu His Cys Arg Ile Lys Thr Thr Gly Ile
 210 215 220

Ser Glu Glu Thr Phe Leu Leu Asn Arg His His Tyr Arg Phe Phe Asp
 225 230 235 240

Val Gly Gly Gln Arg Ser Glu Arg Arg Lys Trp Ile His Cys Phe Glu
 245 250 255

Asn Val Thr Ala Leu Leu Phe Leu Val Ser Leu Ala Gly Tyr Asp Gln
 260 265 270

Cys Leu Val Glu Asp Asn Ser Gly Asn Gln Met Gln Glu Ala Leu Leu
 275 280 285

Leu Trp Asp Ser Ile Cys Asn Ser Ser Trp Phe Ser Glu Ser Ala Met
 290 295 300

Ile Leu Phe Leu Asn Lys Leu Asp Leu Phe Lys Arg Lys Val His Ile
 305 310 315 320

Ser Pro Ile Gln Lys His Phe Pro Asp Tyr Gln Glu Val Gly Ser Thr
 325 330 335

Pro Thr Phe Val Gln Thr Gln Cys Pro Leu Ala Asp Asn Ala Val Arg
 340 345 350

Ser Gly Met Tyr Tyr Phe Tyr Leu Lys Phe Glu Ser Leu Asn Arg Ile
 355 360 365

Ala Ser Arg Ser Cys Tyr Cys His Phe Thr Thr Ala Thr Asp Thr Ser
 370 375 380

Leu Leu Gln Arg Val Met Val Ser Val Gln Asp Thr Ile Met Ser Asn
 385 390 395 400

Asn Leu Glu Phe Asn Leu Val
 405

<210> 11

<211> 407

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: G alpha
transplant

<400> 11

Met	Gly	Cys	Met	Ser	Ser	Lys	Tyr	Ala	Asp	Thr	Ser	Gly	Gly	Glu	Val
1				5					10					15	

Ile	Gln	Lys	Lys	Leu	Ser	Asp	Thr	Gln	Thr	Ser	Asn	Ser	Ser	Thr	Thr
			20					25					30		

Gly	Ser	Gln	Asn	Ala	Arg	Val	Pro	Val	Leu	Glu	Asn	Trp	Leu	Asn	Ile
		35					40					45			

Val	Leu	Arg	Gly	Lys	Pro	Gln	Asn	Val	Glu	Ser	Ser	Gly	Val	Arg	Val
	50					55					60				

Lys	Gly	Asn	Ser	Thr	Ser	Gly	Gly	Asn	Asp	Ile	Lys	Val	Leu	Leu	Leu
65					70					75					80

Gly	Ala	Gly	Asp	Ser	Gly	Lys	Thr	Thr	Ile	Met	Lys	Gln	Met	Arg	Leu
				85					90					95	

Leu	Tyr	Ser	Pro	Gly	Phe	Ser	Gln	Val	Val	Arg	Lys	Gln	Tyr	Arg	Val
			100					105					110		

Met	Ile	Phe	Glu	Asn	Ile	Ile	Ser	Ser	Leu	Cys	Leu	Leu	Leu	Glu	Ala
		115					120				125				

Met	Asp	Asn	Ser	Asn	Val	Ser	Leu	Leu	Pro	Glu	Asn	Glu	Lys	Tyr	Arg
	130					135					140				

Ala	Val	Ile	Leu	Arg	Lys	His	Thr	Ser	Gln	Pro	Asn	Glu	Pro	Phe	Ser
145					150					155					160

Pro	Glu	Ile	Tyr	Glu	Ala	Val	His	Ala	Leu	Thr	Leu	Asp	Thr	Lys	Leu
				165					170					175	

Arg	Thr	Val	Gln	Ser	Cys	Gly	Thr	Asn	Leu	Ser	Leu	Leu	Asp	Asn	Phe
								185						190	

Tyr Tyr Tyr Gln Asp His Ile Asp Arg Ile Phe Asp Pro Gln Tyr Ile
 195 200 205

Pro Ser Asp Gln Asp Ile Leu His Cys Arg Ile Lys Thr Thr Gly Ile
 210 215 220

Ser Glu Glu Thr Phe Leu Leu Asn Arg His His Tyr Arg Phe Phe Asp
 225 230 235 240

Val Gly Gly Gln Arg Ser Glu Arg Arg Lys Trp Ile His Cys Phe Glu
 245 250 255

Asn Val Thr Ala Leu Leu Phe Leu Val Ser Leu Ala Gly Tyr Asp Gln
 260 265 270

Cys Leu Val Glu Asp Asn Ser Gly Asn Gln Met Gln Glu Ala Leu Leu
 275 280 285

Leu Trp Asp Ser Ile Cys Asn Ser Ser Trp Phe Ser Glu Ser Ala Met
 290 295 300

Ile Leu Phe Leu Asn Lys Leu Asp Leu Phe Lys Arg Lys Val His Ile
 305 310 315 320

Ser Pro Ile Gln Lys His Phe Pro Asp Tyr Gln Glu Val Gly Ser Thr
 325 330 335

Pro Thr Phe Val Gln Thr Gln Cys Pro Leu Ala Asp Asn Ala Val Arg
 340 345 350

Ser Gly Met Tyr Tyr Phe Tyr Leu Lys Phe Glu Ser Leu Asn Arg Ile
 355 360 365

Ala Ser Arg Ser Cys Tyr Cys His Phe Thr Thr Ala Thr Asp Thr Ser
 370 375 380

Leu Leu Gln Arg Val Met Val Ser Val Gln Asp Thr Ile Met Ser Asn
 385 390 395 400

Asn Leu Phe Lys Asp Val Arg
 405

<210> 12

<211> 201

<212> PRT

<213> Schizosaccharomyces pombe

<400> 12

Met Lys Ile Thr Ala Val Ile Ala Leu Leu Phe Ser Leu Ala Ala Ala
 1 5 10 15

Ser Pro Ile Pro Val Ala Asp Pro Gly Val Val Ser Val Ser Lys Ser
 20 25 30

Tyr Ala Asp Phe Leu Arg Val Tyr Gln Ser Trp Asn Thr Phe Ala Asn
 35 40 45

Pro Asp Arg Pro Asn Leu Lys Lys Arg Glu Phe Glu Ala Ala Pro Ala
 50 55 60

Lys Thr Tyr Ala Asp Phe Leu Arg Ala Tyr Gln Ser Trp Asn Thr Phe
 65 70 75 80

Val Asn Pro Asp Arg Pro Asn Leu Lys Lys Arg Glu Phe Glu Ala Ala
 85 90 95

Pro Glu Lys Ser Tyr Ala Asp Phe Leu Arg Ala Tyr His Ser Trp Asn
 100 105 110

Thr Phe Val Asn Pro Asp Arg Pro Asn Leu Lys Lys Arg Glu Phe Glu
 115 120 125

Ala Ala Pro Ala Lys Thr Tyr Ala Asp Phe Leu Arg Ala Tyr Gln Ser
 130 135 140

Trp Asn Thr Phe Val Asn Pro Asp Arg Pro Asn Leu Lys Lys Arg Thr
 145 150 155 160

Glu Glu Asp Glu Glu Asn Glu Glu Glu Asp Glu Glu Tyr Tyr Arg Phe
 165 170 175

Leu Gln Phe Tyr Ile Met Thr Val Pro Glu Asn Ser Thr Ile Thr Asp
 180 185 190

Val Asn Ile Thr Ala Lys Phe Glu Ser
 195 200

<210> 13

<211> 99

<212> PRT

<213> Schizosaccharomyces pombe

<400> 13

Met Lys Ile Thr Ala Val Ile Ala Leu Leu Phe Ser Leu Ala Ala Ala
 1 5 10 15

Ser Pro Ile Pro Val Ala Asp Pro Gly Val Val Ser Val Ser Lys Ser
 20 25 30

Tyr Ala Asp Phe Leu Arg Val Tyr Gln Ser Trp Asn Thr Phe Ala Asn
 35 40 45

Pro Asp Arg Pro Asn Leu Lys Lys Arg Thr Glu Glu Asp Glu Glu Asn
 50 55 60

Glu Glu Glu Asp Glu Glu Tyr Tyr Arg Phe Leu Gln Phe Tyr Ile Met
 65 70 75 80

Thr Val Pro Glu Asn Ser Thr Ile Thr Asp Val Asn Ile Thr Ala Lys
 85 90 95

Phe Glu Ser

<210> 14

<211> 415

<212> PRT

<213> Homo sapiens

<400> 14

Met Gly Gly His Pro Gln Leu Arg Leu Val Lys Ala Leu Leu Leu Leu
 1 5 10 15

Gly Leu Asn Pro Val Ser Ala Ser Leu Gln Asp Gln His Cys Glu Ser
 20 25 30

Leu Ser Leu Ala Ser Asn Ile Ser Gly Leu Gln Cys Asn Ala Ser Val
 35 40 45

Asp Leu Ile Gly Thr Cys Trp Pro Arg Ser Pro Ala Gly Gln Leu Val
 50 55 60

Val Arg Pro Cys Pro Ala Phe Phe Tyr Gly Val Arg Tyr Asn Thr Thr
 65 70 75 80

Asn Asn Gly Tyr Arg Glu Cys Leu Ala Asn Gly Ser Trp Ala Ala Arg
 85 90 95

Val	Asn	Tyr	Ser	Glu	Cys	Gln	Glu	Ile	Leu	Asn	Glu	Glu	Lys	Lys	Ser	100	105	110	
Lys	Val	His	Tyr	His	Val	Ala	Val	Ile	Ile	Asn	Tyr	Leu	Gly	His	Cys	115	120	125	
Ile	Ser	Leu	Val	Ala	Leu	Leu	Val	Ala	Phe	Val	Leu	Phe	Leu	Arg	Leu	130	135	140	
Arg	Ser	Ile	Arg	Cys	Leu	Arg	Asn	Ile	Ile	His	Trp	Asn	Leu	Ile	Ser	145	150	155	160
Ala	Phe	Ile	Leu	Arg	Asn	Ala	Thr	Trp	Phe	Val	Val	Gln	Leu	Thr	Met	165	170	175	
Ser	Pro	Glu	Val	His	Gln	Ser	Asn	Val	Gly	Trp	Cys	Arg	Leu	Val	Thr	180	185	190	
Ala	Ala	Tyr	Asn	Tyr	Phe	His	Val	Thr	Asn	Phe	Phe	Trp	Met	Phe	Gly	195	200	205	
Glu	Gly	Cys	Tyr	Leu	His	Thr	Ala	Ile	Val	Leu	Thr	Tyr	Ser	Thr	Asp	210	215	220	
Arg	Leu	Arg	Lys	Trp	Met	Phe	Ile	Cys	Ile	Gly	Trp	Gly	Val	Pro	Phe	225	230	235	240
Pro	Ile	Ile	Val	Ala	Trp	Ala	Ile	Gly	Lys	Leu	Tyr	Tyr	Asp	Asn	Glu	245	250	255	
Lys	Cys	Trp	Phe	Gly	Lys	Arg	Pro	Gly	Val	Tyr	Thr	Asp	Tyr	Ile	Tyr	260	265	270	
Gln	Gly	Pro	Met	Ile	Leu	Val	Leu	Leu	Ile	Asn	Phe	Ile	Phe	Leu	Phe	275	280	285	
Asn	Ile	Val	Arg	Ile	Leu	Met	Thr	Lys	Leu	Arg	Ala	Ser	Thr	Thr	Ser	290	295	300	
Glu	Thr	Ile	Gln	Tyr	Arg	Lys	Ala	Val	Lys	Ala	Thr	Leu	Val	Leu	Leu	305	310	315	320
Pro	Leu	Leu	Gly	Ile	Thr	Tyr	Met	Leu	Phe	Phe	Val	Asn	Pro	Gly	Glu	325	330	335	
Asp	Glu	Val	Ser	Arg	Val	Val	Phe	Ile	Tyr	Phe	Asn	Ser	Phe	Leu	Glu	340	345	350	

Ser Phe Gln Gly Phe Phe Val Ser Val Phe Tyr Cys Phe Leu Asn Ser
 355 360 365

Glu Val Arg Ser Ala Ile Arg Lys Arg Trp His Arg Trp Gln Asp Lys
 370 375 380

His Ser Ile Arg Ala Arg Val Ala Arg Ala Met Ser Ile Pro Thr Ser
 385 390 395 400

Pro Thr Arg Val Ser Phe His Ser Ile Lys Gln Ser Thr Ala Val
 405 410 415

<210> 15

<211> 1224

<212> DNA

<213> Schizosaccharomyces pombe

<400> 15

atgggatgca tgtcgagtaa atacgctgat acatcaggag gagaagtcac tcaaaagaag 60
 ctttcagata cgcaaacctc aaacagctct acaactggaa gtcaaaacgc tcgagttcca 120
 gtccttgaaa actggcttaa tctcgtcctg cgtggaaaac caaaaaatgt ggaaagtctt 180
 ggagtacgcg taaaaggaaa ttctacttca ggtggaaatg acattaaagt tttgctotta 240
 ggcgccggtg atagtgggaa aacgaccatt atgaagcaga tgagattatt gtatagcccc 300
 ggtttttagtc aagtagttag aaagcagtat cgagtgatga tttttgaaaa tatcatctcc 360
 tctctatgtc ttcttcttga agctatggat aatagtaatg tctctttact tccggaaaaat 420
 gagaagtatc gggcagttat cctaagaaaa cacacttctc aaccaaatga gccattttct 480
 ccagaaatat atgaagctgt tcatgccttg acattggata ccaaacttcg tacggtgcaa 540
 agttgtggtg ccaacctctc tttgttagac aatttttatt actatcaaga tcacattgat 600
 cgaatttttg acccacaata tataccttct gatcaagata tccttcaactg tcgtatcaag 660
 acgaccggta tatcagaaga aacattttctg ttaaactcgtc atcattaccg attttttgat 720
 gtaggaggac agagatcaga gcgcagaaaa tggattcatt gctttgaaaa tgtcactgca 780
 ttgtttgttc tcgtttcttt ggcagggttac gatcaatgcc ttgtagagga caattcagga 840
 aatcagatgc aggaggcggt attattatgg gattccatat gtaactctag ctggttttca 900
 gaatcagcaa tgatactttt tctaaataaa cttgatttat ttaaaagaaa gggtcacatt 960
 tccccatcc agaagcattt tcttgattac caagaagttg gttcaacacc aacattcgta 1020
 caaactcaat gccctcttgc cgacaacgca gttcgaagcg gtatgtatta cttttactta 1080
 aagtttgaaa gtcttaaatcg catcgcttct cgtagttgct attgccattt taccacagct 1140
 acagacacta gtttgctcca aagggtaatg gtatccgttc aagatacgat tatgtccaac 1200
 aatctacagt cacttatggt ttag 1224

<210> 16

<211> 1224

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:G alpha
transplant

<400> 16

```
atgggatgca tgtcgagtaa atacgctgat acatcaggag gagaagtcac tcaaaagaag 60
ctttcagata cgcaaacctc aaacagctct acaactggaa gtcaaaacgc tcgagttcca 120
gtccttgaaa actggcttaa tatcgctctg cgtggaaaac cacaaaatgt ggaaagttct 180
ggagtacgcg taaaaggaaa ttctacttca ggtggaaatg acattaaagt tttgctctta 240
ggcgccggtg atagtgggaa aacgaccatt atgaagcaga tgagattatt gtatagcccc 300
ggttttagtc aagtagttag aaagcagtat cgagtgatga tttttgaaaa tatcatctcc 360
tctctatgtc ttcttcttga agctatggat aatagtaatg tctctttact tccggaaaat 420
gagaagtatc gggcagttat cctaagaaaa cacacttctc aaccaatga gccattttct 480
ccagaaatat atgaagctgt tcatgccttg acattggata ccaaacttcg tacggtgcaa 540
agttgtggta ccaacctctc tttgtagac aatttttatt actatcaaga tcacattgat 600
cgaatttttg acccacaata tataccttct gatcaagata tccttactg tcgtatcaag 660
acgaccggta tatcagaaga aacatttctg ttaaactcgtc atcattaccg attttttgat 720
gtaggaggac agagatcaga gcgcagaaaa tggattcatt gctttgaaaa tgtcactgca 780
ttgttgtttc tcgtttcttt ggcagggtac gatcaatgcc ttgtagagga caattcagga 840
aatcagatgc aggaggcggt attattatgg gattccatat gtaactctag ctggttttca 900
gaatcagcaa tgatactttt tctaaataaa cttgatttat ttaaaagaaa gggtcacatt 960
tccccatcc agaagcattt tcctgattac caagaagttg gttcaacacc aacattcgta 1020
caaactcaat gccctcttgc cgacaacgca gttcgaagcg gtatgtatta cttttactta 1080
aagtttgaaa gtcttaaatcg catcgcttct cgtagttgct attgccattt taccacagct 1140
acagacacta gtttgctcca aagggtaatg gtatccgttc aagatacgat tatgtccaac 1200
aatctacaat atgaacttct ttag 1224
```

<210> 17

<211> 1224

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:G alpha
transplant

<400> 17

```
atgggatgca tgtcgagtaa atacgctgat acatcaggag gagaagtcac tcaaaagaag 60
ctttcagata cgcaaacctc aaacagctct acaactggaa gtcaaaacgc tcgagttcca 120
gtccttgaaa actggcttaa tatcgctctg cgtggaaaac cacaaaatgt ggaaagttct 180
ggagtacgcg taaaaggaaa ttctacttca ggtggaaatg acattaaagt tttgctctta 240
ggcgccggtg atagtgggaa aacgaccatt atgaagcaga tgagattatt gtatagcccc 300
ggttttagtc aagtagttag aaagcagtat cgagtgatga tttttgaaaa tatcatctcc 360
tctctatgtc ttcttcttga agctatggat aatagtaatg tctctttact tccggaaaat 420
gagaagtatc gggcagttat cctaagaaaa cacacttctc aaccaatga gccattttct 480
ccagaaatat atgaagctgt tcatgccttg acattggata ccaaacttcg tacggtgcaa 540
agttgtggta ccaacctctc tttgtagac aatttttatt actatcaaga tcacattgat 600
```

cgaatttttg acccacaata tataccttct gatcaagata tccttcaactg tcgtatcaag 660
acgaccggta tatcagaaga aacatttctg ttaaactcgtc atcattaccg attttttgat 720
gtaggaggac agagatcaga gcgcagaaaa tggattcatt gctttgaaaa tgtcaactgca 780
ttgttgtttc tcgtttcttt ggcaggttac gatcaatgcc ttgtagagga caattcagga 840
aatcagatgc aggaggcggtt attattatgg gattccatat gtaactctag ctgggttttca 900
gaatcagcaa tgatactttt tctaaataaaa cttgatttat ttaaaagaaa gggtcacatt 960
tcccccatcc agaagcattt tcctgattac caagaagttg gttcaacacc aacattcgta 1020
caaactcaat gccctcttgc cgacaacgca gttcgaagcg gtatgtatta cttttactta 1080
aagtttgaaa gtcttaatcg catcgcttct cgtagttgct attgccattt taccacagct 1140
acagacacta gtttgctcca aagggtaatg gtatccgttc aagatacgat tatgtccaac 1200
aatctagaat ataactttgt ttag 1224

<210> 18

<211> 1224

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:G alpha
transplant

<400> 18

atgggatgca tgtcgagtaa atacgctgat acatcaggag gagaagtcac tcaaaagaag 60
ctttcagata cgcaaacctc aaacagctct acaactggaa gtcaaaacgc tcgagttcca 120
gtccttgaaa actggcttaa tatcgctcctg cgtggaaaac caaaaaatgt ggaaagttct 180
ggagtacgcg taaaaggaaa ttctacttca ggtggaaatg acattaaagt tttgctctta 240
ggcgccggtg atagtgggaa aacgaccatt atgaagcaga tgagattatt gtatagcccc 300
ggttttagtc aagtagttag aaagcagtat cgagtgatga tttttgaaaa tatcatctcc 360
tctctatgtc ttcttcttga agctatggat aatagtaatg tctctttact tccggaat 420
gagaagtatc gggcagttat octaagaaaa cacacttctc aaccaatga gccattttct 480
ccagaaatat atgaagctgt tcatgccttg acattggata ccaaacttcg tacggtgcaa 540
agttgtggtg ccaacctctc tttgttagac aatttttatt actatcaaga tcacattgat 600
cgaatttttg acccacaata tataccttct gatcaagata tccttcaactg tcgtatcaag 660
acgaccggta tatcagaaga aacatttctg ttaaactcgtc atcattaccg attttttgat 720
gtaggaggac agagatcaga gcgcagaaaa tggattcatt gctttgaaaa tgtcaactgca 780
ttgttgtttc tcgtttcttt ggcaggttac gatcaatgcc ttgtagagga caattcagga 840
aatcagatgc aggaggcggtt attattatgg gattccatat gtaactctag ctgggttttca 900
gaatcagcaa tgatactttt tctaaataaaa cttgatttat ttaaaagaaa gggtcacatt 960
tcccccatcc agaagcattt tcctgattac caagaagttg gttcaacacc aacattcgta 1020
caaactcaat gccctcttgc cgacaacgca gttcgaagcg gtatgtatta cttttactta 1080
aagtttgaaa gtcttaatcg catcgcttct cgtagttgct attgccattt taccacagct 1140
acagacacta gtttgctcca aagggtaatg gtatccgttc aagatacgat tatgtccaac 1200
aatctaggat gcggacttta ttag 1224

<210> 19

<211> 1224

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:G alpha
transplant

<400> 19

```
atgggatgca tgtcgagtaa atacgctgat acatcaggag gagaagtcac tcaaaagaag 60
ctttcagata cgcaaaccctc aaacagctct acaactggaa gtcaaaacgc tcgagttcca 120
gtccttgaaa actggcttaa tatcgctcctg cgtggaaaac cacaaaatgt ggaaagttct 180
ggagtacgcg taaaaggaaa ttctacttca ggtggaaatg acattaaagt tttgctctta 240
ggcgccggtg atagtgggaa aacgaccatt atgaagcaga tgagattatt gtatagcccc 300
ggtttttagtc aagtagtttag aaagcagtat cgagtgatga tttttgaaaa tatcatctcc 360
tctctatgtc ttcttcttga agctatggat aatagtaatg tctctttact tccggaaaat 420
gagaagtatc gggcagttat cctaagaaaa cacacttctc aaccaatga gccattttct 480
ccagaaatat atgaagctgt tcatgccttg acattggata ccaaacttcg tacggtgcaa 540
agttgtggtg ccaacctctc ttgttttagac aatttttatt actatcaaga tcacattgat 600
cgaatttttg acccacaata tataccttct gatcaagata tccttctact tcgtatcaag 660
acgaccggtg tatcagaaga aacattttctg ttaaactcgtc atcattaccg attttttgat 720
gtaggaggac agagatcaga gcgcagaaaa tggattcatt gctttgaaaa tgtcactgca 780
ttgttgtttc tcgtttcttt ggcagggttac gatcaatgcc ttgtagagga caattcagga 840
aatcagatgc aggaggcggt attattatgg gattccatat gtaactctag ctgggtttca 900
gaatcagcaa tgatactttt tctaaataaaa cttgatttat ttaaaagaaa gggttcacatt 960
tcccccatcc agaagcattt tcttgattac caagaagtgt gttcaacacc aacattcgtg 1020
caaactcaat gccctcttgc cgacaacgca gttcgaagcg gtatgtatta cttttactta 1080
aagtttgaaa gtcttaaatcg catcgcttct cgtagttgct attgccattt taccacagct 1140
acagacacta gtttgctcca aagggtaatg gtatccgttc aagatacgat tatgtccaac 1200
aatctagatt ggggactttt ttag                                     1224
```

<210> 20

<211> 1224

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:G alpha
transplant

<400> 20

```
atgggatgca tgtcgagtaa atacgctgat acatcaggag gagaagtcac tcaaaagaag 60
ctttcagata cgcaaaccctc aaacagctct acaactggaa gtcaaaacgc tcgagttcca 120
gtccttgaaa actggcttaa tatcgctcctg cgtggaaaac cacaaaatgt ggaaagttct 180
ggagtacgcg taaaaggaaa ttctacttca ggtggaaatg acattaaagt tttgctctta 240
ggcgccggtg atagtgggaa aacgaccatt atgaagcaga tgagattatt gtatagcccc 300
ggtttttagtc aagtagtttag aaagcagtat cgagtgatga tttttgaaaa tatcatctcc 360
tctctatgtc ttcttcttga agctatggat aatagtaatg tctctttact tccggaaaat 420
```

```

gagaagtatc gggcagttat cctaagaaaa cacacttctc aacccaatga gccattttct 480
ccagaaatat atgaagctgt tcatgccttg acattggata ccaaacttcg tacggtgcaa 540
agttgtggta ccaacctctc tttgttagac aatttttatt actatcaaga tcacattgat 600
cgaatttttg acccacaata tataccttct gatcaagata tccttcaactg tcgtatcaag 660
acgaccggta tatcagaaga aacatttctg ttaaactcgtc atcattaccg attttttgat 720
gtaggaggac agagatcaga gcgcagaaaa tggattcatt gctttgaaaa tgtcactgca 780
ttgttgtttc tcgtttcttt ggcaggttac gatcaatgcc ttgtagagga caattcagga 840
aatcagatgc aggaggcgtt attattatgg gattccatat gtaactctag ctgggttttca 900
gaatcagcaa tgatactttt tctaaataaa cttgatttat taaaagaaa gggttcacatt 960
tcccccatcc agaagcattt tcctgattac caagaagttg gttcaacacc aacattcgta 1020
caaactcaat gccctcttgc cgacaacgca gttcgaagcg gtatgtatta cttttactta 1080
aagtttgaaa gtcttaatcg catcgcttct cgtagttgct attgccattt taccacagct 1140
acagacacta gtttgctcca aagggtaatg gtatccgttc aagatacgat tatgtccaac 1200
aatctagaat gcggacttta ttag
1224

```

<210> 21

<211> 1224

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:G alpha
transplant

<400> 21

```

atgggatgca tgtcgagtaa atacgctgat acatcaggag gagaagtcac tcaaaagaag 60
ctttcagata cgcaaacctc aaacagctct acaactggaa gtcaaaacgc tcgagttcca 120
gtccttgaaa actggcttaa tctcgtcttg cgtggaaaac cacaaaatgt ggaaagttct 180
ggagtacgcg taaaaggaaa ttctacttca ggtggaaatg acattaaagt tttgctctta 240
ggcgccgggtg atagtgggaa aacgaccatt atgaagcaga tgagattatt gtatagcccc 300
ggttttagtc aagtagttag aaagcagtat cgagtgatga ttttgaaaa tatcatctcc 360
tctctatgtc ttcttcttga agctatggat aatagtaatg tctctttact tccggaaaat 420
gagaagtatc gggcagttat cctaagaaaa cacacttctc aacccaatga gccattttct 480
ccagaaatat atgaagctgt tcatgccttg acattggata ccaaacttcg tacggtgcaa 540
agttgtggta ccaacctctc tttgttagac aatttttatt actatcaaga tcacattgat 600
cgaatttttg acccacaata tataccttct gatcaagata tccttcaactg tcgtatcaag 660
acgaccggta tatcagaaga aacatttctg ttaaactcgtc atcattaccg attttttgat 720
gtaggaggac agagatcaga gcgcagaaaa tggattcatt gctttgaaaa tgtcactgca 780
ttgttgtttc tcgtttcttt ggcaggttac gatcaatgcc ttgtagagga caattcagga 840
aatcagatgc aggaggcgtt attattatgg gattccatat gtaactctag ctgggttttca 900
gaatcagcaa tgatactttt tctaaataaa cttgatttat taaaagaaa gggttcacatt 960
tcccccatcc agaagcattt tcctgattac caagaagttg gttcaacacc aacattcgta 1020
caaactcaat gccctcttgc cgacaacgca gttcgaagcg gtatgtatta cttttactta 1080
aagtttgaaa gtcttaatcg catcgcttct cgtagttgct attgccattt taccacagct 1140
acagacacta gtttgctcca aagggtaatg gtatccgttc aagatacgat tatgtccaac 1200
aatctatata ttggactttg cttag
1224

```

<210> 22
<211> 1224
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:G alpha
transplant

<400> 22
atgggatgca tgtcgagtaa atacgctgat acatcaggag gagaagtcac tcaaaagaag 60
ctttcagata cgcaaacctc aaacagctct acaactggaa gtcaaaacgc tcgagttcca 120
gtccttgaaa actggcttaa tategtcttg cgtggaaaac cacaaaatgt ggaaagttct 180
ggagtacgcg taaaaggaaa ttctacttca ggtggaaatg acattaaagt tttgctctta 240
ggcgccggtg atagtgggaa aacgaccatt atgaagcaga tgagattatt gtatagcccc 300
ggttttagtc aagtagttag aaagcagtat cgagtgatga tttttgaaa tatcatctcc 360
tctctatgtc ttcttcttga agctatggat aatagtaatg tctctttact tccggaaaat 420
gagaagtatc gggcagttat cctaagaaaa cacacttctc aacccaatga gccatcttct 480
ccagaaatat atgaagctgt tcatgccttg acattggata ccaaacttcg tacggtgcaa 540
agttgtggta ccaacctctc tttgtagac aatTTTTtatt actatcaaga tcacattgat 600
cgaatttttg acccacaata tataccttct gatcaagata tctttcactg tcgtatcaag 660
acgaccggta tatcagaaga aacatttctg ttaaatcgtc atcattaccg atTTTTtgat 720
gtaggaggac agagatcaga gcgcagaaaa tggattcatt gctttgaaaa tgtcactgca 780
ttgttgtttc tcgtttcttt ggcaggttac gatcaatgcc ttgtagagga caattcagga 840
aatcagatgc aggaggcgtt attattatgg gattccatat gtaactctag ctgggtttca 900
gaatcagcaa tgatactttt tctaaataaa cttgatttat ttaaaagaaa gggtcacatt 960
tccccatcc agaagcattt tctgattac caagaagttg gttcaacacc aacattcgta 1020
caaactcaat gccctcttgc cgacaacgca gttcgaagcg gtatgtatta cttttactta 1080
aagtttgaaa gtcttaatcg catcgcttct cgtagttgct attgccattt taccacagct 1140
acagacacta gtttgctcca aagggtaatg gtatccgttc aagatacgat tatgtccaac 1200
aatctagata ttatgcttca atag 1224

<210> 23
<211> 1224
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:G alpha
transplant

<400> 23
atgggatgca tgtcgagtaa atacgctgat acatcaggag gagaagtcac tcaaaagaag 60
ctttcagata cgcaaacctc aaacagctct acaactggaa gtcaaaacgc tcgagttcca 120
gtccttgaaa actggcttaa tategtcttg cgtggaaaac cacaaaatgt ggaaagttct 180
ggagtacgcg taaaaggaaa ttctacttca ggtggaaatg acattaaagt tttgctctta 240

ggcgccggtg atagtgggaa aacgaccatt atgaagcaga tgagattatt gtatagcccc 300
 ggttttagtc aagtagttag aaagcagtat cgagtgatga tttttgaaaa tatcatctcc 360
 tctctatgtc ttcttcttga agctatggat aatagtaatg tctctttact tccggaaaaat 420
 gagaagtatc gggcagttat cctaagaaaa cacacttctc aaccaatga gccattttct 480
 ccagaaatat atgaagctgt tcatgccttg acattggata ccaaacttcg tacggtgcaa 540
 agttgtggta ccaacctctc ttgttagac aatttttatt actatcaaga tcacattgat 600
 cgaatttttg acccacaata tataccttct gatcaagata tccttcaactg tcgtatcaag 660
 acgaccggta tatcagaaga aacatttctg ttaaactcgtc atcattaccg attttttgat 720
 gtaggaggac agagatcaga gcgcagaaaa tggattcatt gctttgaaaa tgtaactgca 780
 ttgttgttctc tcgtttcttt ggcaggttac gatcaatgcc ttgtagagga caattcagga 840
 aatcagatgc aggaggcgtt attattatgg gattccatat gtaactctag ctggttttca 900
 gaatcagcaa tgatactttt tctaaataaa cttgatttat ttaaaagaaa ggttcacatt 960
 tcccccatcc agaagcattt tcctgattac caagaagttg gttcaacacc aacattcgta 1020
 caaactcaat gccctcttgc cgacaacgca gttcgaagcg gtatgtatta cttttactta 1080
 aagtttgaaa gtcttaatcg catcgcttct cgtagttgct attgccattt taccacagct 1140
 acagacacta gtttgctcca aagggtaatg gtatccgttc aagatacgat tatgtccaac 1200
 aatctacaac ttatgcttca atag 1224

<210> 24

<211> 1224

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:G alpha
transplant

<400> 24

atgggatgca tgcagagtaa atacgtgat acatcaggag gagaagtcac tcaaaagaag 60
 ctttcagata cgaaacctc aaacagctct acaactggaa gtcaaaacgc tcgagttcca 120
 gtccttgaaa actggcttaa tatcgtcctg cgtggaaaac cacaaaatgt ggaaagttct 180
 ggagtacgcg taaaaggaaa ttctacttca ggtggaaatg acattaaagt ttgctctta 240
 ggcgccggtg atagtgggaa aacgaccatt atgaagcaga tgagattatt gtatagcccc 300
 ggttttagtc aagtagttag aaagcagtat cgagtgatga tttttgaaaa tatcatctcc 360
 tctctatgtc ttcttcttga agctatggat aatagtaatg tctctttact tccggaaaaat 420
 gagaagtatc gggcagttat cctaagaaaa cacacttctc aaccaatga gccattttct 480
 ccagaaatat atgaagctgt tcatgccttg acattggata ccaaacttcg tacggtgcaa 540
 agttgtggta ccaacctctc ttgttagac aatttttatt actatcaaga tcacattgat 600
 cgaatttttg acccacaata tataccttct gatcaagata tccttcaactg tcgtatcaag 660
 acgaccggta tatcagaaga aacatttctg ttaaactcgtc atcattaccg attttttgat 720
 gtaggaggac agagatcaga gcgcagaaaa tggattcatt gctttgaaaa tgtaactgca 780
 ttgttgttctc tcgtttcttt ggcaggttac gatcaatgcc ttgtagagga caattcagga 840
 aatcagatgc aggaggcgtt attattatgg gattccatat gtaactctag ctggttttca 900
 gaatcagcaa tgatactttt tctaaataaa cttgatttat ttaaaagaaa ggttcacatt 960
 tcccccatcc agaagcattt tcctgattac caagaagttg gttcaacacc aacattcgta 1020
 caaactcaat gccctcttgc cgacaacgca gttcgaagcg gtatgtatta cttttactta 1080
 aagtttgaaa gtcttaatcg catcgcttct cgtagttgct attgccattt taccacagct 1140

```

acagacacta gtttgctcca aagggtaatg gtatccgttc aagatacgat tatgtccaac 1200
aatctagaat ttaatcttgt ttag                                     1224

```

<210> 25

<211> 1224

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:G alpha
transplant

<400> 25

```

atgggatgca tgctgagtaa atacgctgat acatcaggag gagaagtcac tcaaaagaag 60
ctttcagata cgcaaacctc aaacagctct acaactggaa gtcaaaacgc tcgagttcca 120
gtccttgaaa actggcttaa tatcgctctg cgtggaaaac cacaaaatgt ggaaagttct 180
ggagtacgcg taaaaggaaa ttctacttca ggtggaaatg acattaaagt tttgctctta 240
ggcgccggtg atagtgggaa aacgaccatt atgaagcaga tgagattatt gtatagcccc 300
ggttttagtc aagtagttag aaagcagtat cgagtgatga tttttgaaaa tatcatctcc 360
tctctatgtc ttcttcttga agctatggat aatagtaatg tctctttact tccggaaaat 420
gagaagtatc gggcagttat cctaagaaaa cacacttctc aaccaatga gccattttct 480
ccagaaaatat atgaagctgt tcatgccttg acattggata ccaaacttcg tacggtgcaa 540
agttgtggta ccaacctctc tttgttagac aatttttatt actatcaaga tcacattgat 600
cgaatttttg acccacaata tataccttct gatcaagata tccttctact tcgtatcaag 660
acgaccggta tatcagaaga aacatttctg ttaaactcgc atcattaccg attttttgat 720
gtaggaggac agagatcaga gcgcagaaaa tggattcatt gctttgaaaa tgctactgca 780
ttgttggttc tcgtttcttt ggcaggttac gatcaatgcc ttgtagagga caattcagga 840
aatcagatgc aggaggcggt attattatgg gattccatat gtaactctag ctggttttca 900
gaatcagcaa tgatactttt tctaaataaa cttgatttat ttaaaagaaa ggttcacatt 960
tccccatcc agaagcattt tcctgattac caagaagttg gttcaacacc aacattcgta 1020
caaaactcaat gccctcttgc cgacaacgca gttcgaagcg gtatgtatta cttttactta 1080
aagtttgaaa gtcttaaatcg catcgcttct cgtagttgct attgccattt taccacagct 1140
acagacacta gtttgctcca aagggtaatg gtatccgttc aagatacgat tatgtccaac 1200
aatctagaaa ttaatcttct ttag                                     1224

```

<210> 26

<211> 625

<212> DNA

<213> Schizosaccharomyces pombe

<400> 26

```

ggggatccac catgaagatc accgctgtca ttgccctttt attctcaatt gctgctgcct 60
cacctattcc agttgcgat cctgggtgtg tttcagttag caagtcatat gctgatttcc 120
ttcgtgttta ccaaagttgg aacacttttg ctaatcctga tagaccaaac ttgaaaaagc 180
gcgaattcga agctgctccc gcaaaaactt atgctgattt ccttcgtgct tatcaaagtt 240
ggaacacttt tgtaaatcct gacagacca atttgaaaaa gcgtgagttt gaagctgccc 300

```

cagagaagag ttatgctgat ttccttcgtg cttaccatag ttggaacact tttgttaatc 360
 ctgacagacc caacttgaaa aagcgcgaat tcgaagctgc tcccgcacaaa acttatgctg 420
 atttccttcg tgcttaccaa agttggaaca cttttgttaa tcctgacaga cccaacttga 480
 aaaagcgcac tgaagaagat gaagagaatg aggaagagga tgaagaatac tatcgctttc 540
 ttcagtttta tatcatgact gtcccagaga attccactat tacagatgtc aatattactg 600
 ccaaatttga gagctaagga tcccc 625

<210> 27

<211> 319

<212> DNA

<213> Schizosaccharomyces pombe

<400> 27

ggggatccac catgaagatc accgctgtca ttgccctttt attctcaatt gctgctgcct 60
 cacctattcc agttgccgat cctggtgtgg tttcagttag caagtcatat gctgatttcc 120
 ttcgtgttta ccaaagttgg aacacttttg ctaatcctga tagacccaac ttgaaaaagc 180
 gactgaaga agatgaagag aatgaggaag aggatgaaga atactatcgc tttcttcagt 240
 tttatatcat gactgtccca gagaattcca ctattacaga tgtcaatatt actgccaaat 300
 ttgagagcta aggatcccc 319

<210> 28

<211> 1248

<212> DNA

<213> Homo sapiens

<400> 28

atgggagggc accgcagct ccgtctcgtc aaggcccttc tccttctggg gctgaacccc 60
 gtctctgcct ccctccagga ccagcactgc gagagcctgt ccctggccag caacatctca 120
 ggactgcagt gcaacgcata cgtggacctc attggcacct gctggccccg cagccctgcg 180
 gggcagctag tggttcggcc ctgccctgcc tttttctatg gtgtccgcta caataccaca 240
 aacaatggct accgggagtg cctggccaat ggcagctggg ccgcccgcgt gaattactcc 300
 gagtgccagg agatcctcaa tgaggagaaa aaaagcaagg tgactacca tgtcgcagtc 360
 atcatcaact acctgggcca ctgtatctcc ctggtggccc tcctggtggc ctttgtctcc 420
 tttctgcggc tcaggagcat ccggtgcctg cgaaacatca tccactggaa cctcatctcc 480
 gccttcatcc tgcgcaacgc cacctggttc gtggtccagc taaccatgag ccccgaggtc 540
 caccagagca acgtgggctg gtgcagggtg gtgacagccg cctacaacta cttccatgtg 600
 accaacttct tctggatgtt cggcgagggc tgetacctgc acacagccat cgtgctcacc 660
 tactccactg accggtgcg caaatggatg ttcactctga ttggctgggg tgtgcccttc 720
 cccatcattg tggcctgggc cattgggaag ctgtactacg acaatgagaa gtgctggttt 780
 ggcaaaaggc ctggggtgta caccgactac atctaccagg gcccatgat cctggctctg 840
 ctgatcaatt tcatcttctt tttcaacatc gtccgcatac tcatgaccaa gctccgggca 900
 tccaccacgt ctgagaccat tcagtacagg aaggctgtga aagccactct ggtgctgctg 960
 cccctcctgg gcatcaccta catgctgttc ttcgtcaatc ccggggagga tgaggtctcc 1020
 cgggtcgtct tcatctactt caactccttc ctggaatcct tccagggtct ctttgtgtct 1080
 gtgttctact gtttctcaa tagtgaggtc cgttctgcc tccggaagag gtggcaccgg 1140
 tggcaggaca agcactcgat ccgtgcccca gtggcccggt ccatgtccat cccacctcc 1200

ccaacccgtg tcagctttca cagcatcaag cagtccacag cagtctga

1248

<210> 29

<211> 8800

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:pREP3Xr

<400> 29

```
aagcttgcac gcctgcaggt cgatcgactc tagaggatca gaaaattatc gccataaaaag 60
acagaataag tcatcagcgg ttgtttcatt tectatatatt tttttttatt tttttatttt 120
ttaataaggg aaaatttaac gtctaaggat acagaagatt gtagcacat taaagtaata 180
aaggcttaag tagtaagtgc cttagcatgt tattgtattt caaaggacat aatctaaaaat 240
aataacaata tcattttctca caagttattc aattttcttt tttttttcta ataatatcaa 300
gaatgtatta tttgtttgac ataagtcaac taattttatt aatatgctgg attaatcttg 360
cagacatgta aattaacaag ttttagtcaa ataacgttga agtttcaatg aactcaaata 420
atttctcttt ttttttatat aaccatatgt ctaatctgat ttatatattt cgcagggatc 480
aactgaagtt atgacatttg gattggatca cttataacct tggtcgcaa ataatacaaa 540
aatcagcgtt ataaaacaaa gaagggtttt gttaagaaat taatcctctt tcttgataag 600
aaagttgaac cgaaattgca gatactgata tatgaaaata ataccacaaa ttttggaat 660
agcgcaagcc tcaatttaaa caataggtga ggacacatga taatgacctc aatgattggt 720
agaagaaaag agcctcatta caaatcgaa aaatgaatgg ttgggtacaa gtttccaaaa 780
catggtaaag tggactttgc gtatgagacg taaatagaaa aaaacacttg ttatatgttt 840
tctagaatta ttgttgtctc tttatgggtg gatgatgcaa aatagtaatt tcggttagtt 900
gctgtaaaac accacgagac aaatagatat ggatatttat taaatcagga aaaacgtaac 960
tctcggctac tggatggttc agtcacccaa cgattactgg ggagagaaaa cagggcaaaa 1020
gcaaagctta aaggaatccg attgtcattc ggcaatgtgc agcgaaacta aaaaccggat 1080
aatggacctg ttaatcgaaa cattgaagat atataaagga agagggaatcc tggcatatca 1140
tcaattgaat aagttgaatt aattatttca atctcattct cactttctga cttatagtcg 1200
ctttgttaaa ttggcctcga ggtcgactct agagatatcg gatccccggg taaaaggaat 1260
gtctcccttg ccagtactgc taggggtttt ctttcaaact atggaagccc attcaagctg 1320
catattacga tttgtttttt cgctttttaga aagtggttta gatgagataa tagaaaaatt 1380
cttgatctcc gacaacgagt acttttatatt tttttgctaa tcactttact caatattagc 1440
tcgaaatcgt agaaacgtag acgggtgcgg gataccgagt ggtgtagtta agaattttta 1500
taaaccacgt ggcccaaaaa tatgaaccca aaacgtttat acatgagtat actttaagaa 1560
ggctataccc ctctgtgtta gatgtagttt tagctaccca acccgagtct atgagcttga 1620
cttcagatgt agaaggcatt aaatcgtttt gaatattaat taaaaaacga tgaaaattaa 1680
atatttaaaa gcaatcatat gctgaaaatt tagtgctgtg gctaatcctt caacatggaa 1740
atgccataaa agtgactttg acaaaaaaaa aagtatatac aggtagtaaa ctcatctact 1800
tcattgactt tgtttacagc atgtggaagg aggaatattt attgctaaat cgtagtttaa 1860
cattcaataa gtaatactat tgaaattcga caagattggc cgcattggatg aaaaagaggc 1920
attttgcttt gggagaatta gttcaaatta gaactgaaaa aaaaaacttt acgaggcaaa 1980
aatgtcggat tgagatcgta aaagttcgct cgctcgtctt tgctttgtga ttgttttcat 2040
ggatacatct tgctggatat ttaaatttta gtactatgta taagatattc tataaatgtt 2100
ttatcaccca aacctgttag cgccttctta attctattca atctggcttt tgctctgaga 2160
```

ctacttcttg gactttcact acttggttagt tatacgggaat ttgtgtaatt agaagtgaaa 2220
taatcctttc tattagtaat gcgagctcga attcgagtct aactccttaa ccactcggac 2280
atagtgaact atctgacact cattgtaaat taatatatat ataggcattt tgttttagtta 2340
aaggtaacta agtaattagt ataaacgaac caattttata atcaggaagt taagtgaatg 2400
gtagcacatg tcgtaaaaaat tgtgaatttt tattgaataa tattttaaat acaagccttt 2460
ctagactagg tatactcata aacatatatg agcaaaagga tagaggagat tacattgcat 2520
cttctacaaa tttattttatt gccctttact gaaaaattaa ataatgagta ctaaatagata 2580
aaaagcgctc agtacaagaa agattgcaaa aatattgcat tcttcatgaa ttaaagttgc 2640
atataaggca tattgaaagt aatagtacta aaacagcagt tagcgaaaat taatagaatt 2700
atattcgcaa gacaattgtg acattaaatt aaaaaattgt agaattttta ctatcctctt 2760
taacgccatg agcctttata aaaaggttaa attagtttta acattctttt tttgagtaag 2820
agtttacaat ttatcaaaac ctgtgttatt atattcatta gtttcaattt attagcatct 2880
agagaaaaat caattggcag ttacattgtt ggaatttatg aagaaaaaga atctacaacg 2940
gagaataagt tgctgatcgc tttccctaaa attgtatatatt ttgctgagct tattttgaca 3000
tttcggtgaa gttttctcta attcgcatc attttaagta aaacaatgag aaataaaaatt 3060
acaaaaata caaattaaaa tacaattttt agctataatt atagacgatg cccttgatc 3120
ccattctgtc tcgcttgccc ctacttttta tcttttatat accataatga acgctgccgc 3180
tactaaccat acccgcattt tacatttcgg actcccaagg acgtacaaaa tagaaaacta 3240
tagaaaaaaa taatcagaaa atagcatgct atctctttgt aaaacgcgtt tgcaagaaga 3300
aaggaaacaa tggagaagag atcatccatt tgtatgtaaa ttttagtaaa cttgaagaaa 3360
tcactaaciaa cttctcttac ttagggattc tatgcaaaac cttgtaaatc atctgatgga 3420
ggactcgatt taatgaattg gaaggttga attcactggc cgctggttta caacgtcgtg 3480
actgggaaaa ccctggcggt acccaactta atcgccctgc agcacatccc cctttcgcca 3540
gctggcgtaa tagcgaagag gccgcaccg atcgccctc ccaacagttg cgcagcctga 3600
atggcgaatg gcgcctgat cggtattttc tccttacgca tctgtgcggt atttcacacc 3660
gcatacgtca aagcaaccat agtacgcgcc ctgtagcggc gcattaagcg cggcgggtgt 3720
ggtggttacg cgcagcgtga ccgctacact tgccagcgcc tttagcggcg ctcctttcgc 3780
ttcttccct tcctttctcg ccacgttcgc cggctttccc cgtcaagctc taaatcgggg 3840
gctcccttta gggttccgat ttagtgcttt acggcacctc gaccccaaaa aacttgattt 3900
gggtgatggt tcacgtagt ggacatcgcc ctgatagacg gtttttcgcc ctttgacgtt 3960
ggagtccacg ttctttaata gtggactctt gttccaaact ggaacaacac tcaactctat 4020
ctcgggctat tcttttgatt tataagggat ttgcccatt tcggtctatt gggttaaaaa 4080
tgagctgatt taacaaaaat ttaacgcgaa ttttaacaaa atattaacgt ttacaatttt 4140
atggtgcact ctcagtacaa tctgctctga tgccgcatag ttaagccagc cccgacaccc 4200
gccaacaccc gctgacgcgc cctgacgggc ttgtctgctc ccggcatccg cttacagaca 4260
agctgtgacc gtctccggga gctgcatgtg tcagaggttt tcaccgtcat caccgaaacg 4320
cgcgagacga aagggcctcg tgatacgcct atttttatag gttaatgtca tgataataat 4380
ggtttcttag acgtcaggtg gcacttttcg gggaaatgtg cgcggaaccc ctatttgttt 4440
atttttctaa atacattcaa atatgtatcc gctcatgaga caataaccct gataaatgct 4500
tcaataatat tgaaaaagga agagtatgag tattcaacat ttccgtgtcg cccttattcc 4560
cttttttgcg gcattttgccc ttctgtttt tgctcaccca gaaacgctgg tgaaagtaaa 4620
agatgctgaa gatcagttgg gtgcacgagt gggttacatc gaactggatc tcaacagcgg 4680
taagatcctt gagagttttc gccccgaaga acgttttcca atgatgagca cttttaaagt 4740
tctgctatgt ggcgcggtat tatcccgtat tgacgccggg caagagcaac tcggtcgccg 4800
catacactat tctcagaatg acttggttga gtactacca gtcacagaaa agcatcttac 4860
ggatggcatg acagtaagag aattatgcag tgctgccata accatgagtg ataacactgc 4920
ggccaactta cttctgacaa cgatcggagg accgaaggag ctaaccgctt ttttgacaaa 4980
catgggggat catgtaactc gccttgatcg ttgggaaccg gagctgaatg aagccatacc 5040

aaacgacgag cgtgacacca cgatgcctgt agcaatggca acaacgttgc gcaaactatt 5100
aactggcgaa ctacttactc tagcttcccc gcaacaatta atagactgga tggaggcgga 5160
taaagttgca ggaccacttc tgcgctcggc ccttccggct ggctggttta ttgctgataa 5220
atctggagcc ggtgagcgtg ggtctcgcgg tatcattgca gcaactgggc cagatggtaa 5280
gccctcccgt atcgtagtta tctacacgac ggggagtcag gcaactatgg atgaacgaaa 5340
tagacagatc gctgagatag gtgcctcact gattaagcat tggtaactgt cagaccaagt 5400
ttactcatat atacttttaga ttgattttaa acttcatttt taatttaaaa ggatctaggt 5460
gaagatcctt tttgataatc tcatgaccaa aatcccttaa cgtgagtttt cgttccactg 5520
agcgtcagac cccgtagaaa agatcaaagg atcttcttga gatccttttt ttctgcgcgt 5580
aatctgctgc ttgcaaacaa aaaaaccacc gctaccagcg gtggttttgt tgcgggatca 5640
agagctacca actctttttc cgaaggtaac tggcttcagc agagcgcaga taccaaatac 5700
tgtccttcta gtgtagccgt agttaggcca ccacttcaag aactctgtag caccgcctac 5760
atacctcgct ctgctaatac tgttaccagt ggctgctgcc agtggcgata agtcgtgtct 5820
taccgggttg gactcaagac gatagttacc ggataaggcg cagcggtcgg gctgaacggg 5880
gggttcgtgc acacagccca gcttggagcg aacgacctac accgaactga gatactaca 5940
gcgtgagcta tgagaaagcg ccacgcttcc cgaagggaga aaggcggaca ggtatccgg 6000
aagcggcagg gtcggaacag gagagcgcac gagggagctt ccagggggaa acgcctggta 6060
tctttatagt cctgtcgggt ttgccacct ctgacttgag cgtcgatttt tgtgatgtc 6120
gtcagggggg cggagcctat ggaaaaacgc cagcaacgcg gcctttttac ggttcctggc 6180
cttttgctgg ccttttgctc acatgttctt tctgcgtta tcccctgatt ctgtggataa 6240
ccgtattacc gcctttgagt gagctgatac cgctcgcgcg agccgaacga ccgagcgcag 6300
cgagtcagt agcgaggaag cggaaagagcg cccaatacgc aaaccgcctc tccccgcgcg 6360
ttggccgatt cattaatgca gctggcacga caggtttccc gactggaaag cgggcagtga 6420
gcgcaacgca attaatgtga gttagctcac tcattaggca cccaggctt tacactttat 6480
gcttccggct cgtatgttgt gtggaattgt gagcgataa caatttcaca caggaaacag 6540
ctatgaccat gattacgcca agcttgcga tcgactacgt cgtaaaggcc gtttctgaca 6600
gagtaaaatt cttgagggaa ctttcacat tatgggaaat ggttcaagaa ggtattgact 6660
taaactccat caaatggtca ggtcattgag tgttttttat ttgttgatt ttttttttt 6720
tagagaaaat cctccaatat caaattagga atcgtagttt catgattttc tgttacacct 6780
aactttttgt gtggtgccct cctccttgct aatattaatg ttaaagtga attctttttc 6840
cttatcacgt tgagccatta gtatcaattt gcttacctgt attcctttac tatcctcctt 6900
tttctccttc ttgataaatg tatgtagatt gcgtatatag tttcgtctac cctatgaaca 6960
tattccattt tgtaatttcg tgcgtttct attatgaatt tcatttataa agtttatgta 7020
caaatatcat aaaaaaagag aatcttttta agcaaggatt ttcttaactt cttcggcgac 7080
agcatcaccg acttcgggtg tactgttgga accacctaaa tcaccagttc tgatacctgc 7140
atccaaaacc tttttaactg catcttcaat ggccttacct tcttcaggca agttcaatga 7200
caatttcaac atcattgcag cagacaagat agtggcgata gggtaacct tattctttgg 7260
caaactgga gcagaaccgt ggcattgttc gtacaaacca aatgcgggtg tcttgctctg 7320
caaagaggcc aaggacgcag atggcaacaa acccaaggaa cctgggataa cggaggcttc 7380
atcggagatg atgtcaccaa acatgttgct ggtgattata ataccattta ggtgggttg 7440
gttcttaact aggatcatgg cggcagaatc aatcaattga tgttgaacct tcaatgtagg 7500
gaattcgttc ttgatggttt cctccacagt ttttctccat aatcttgaag aggccaaaac 7560
attagcttta tccaaggacc aaataggcaa tgggtggctca tgttgtaggg ccatgaaagc 7620
ggccattctt gtgattcttt gcacttctgg aacggtgtat tgttcactat cccaagcgac 7680
accatcacca tctcttctt tctctttacc aaagtaaata cctcccacta attctctgac 7740
aacaacgaag tcagtacct tagcaaatg tggcttgatt ggagataagt ctaaaagaga 7800
gtcggatgca aagttacatg gtcttaagtt ggcgtacaat tgaagttctt tacggatttt 7860
tagtaaacct tgttcaggtc taacactacc ggtaccccat ttaggaccac ccacagcacc 7920

taacaaaacg gcatcaacct tcttggagggc ttccagcgcc tcatctggaa gtgggacacc 7980
tgtagcatcg atagcagcac caccaattaa atgattttcg aaatcgaact tgacattgga 8040
acgaacatca gaaatagctt taagaacctt aatggcttcg gctgtgattt cttgaccaac 8100
gtggtcacct ggcaaaacga cgatcttctt aggggcagac ataggggcag acattagaat 8160
ggtatatcct tgaaatatat atatatttg ctgaaatgta aaaggtaaga aaagttagaa 8220
agtaagacga ttgctaacca cctattggaa aaaacaatag gtccttaaata aatattgtca 8280
acttcaagta ttgtgatgca agcatttagt catgaacgct tctctattct atatgaaaag 8340
ccggttccgg cctctcacct ttcccttttc tcccaatttt tcagttgaaa aaggatatatg 8400
cgtcaggcga cctctgaaat taacaaaaaa ttccagtcga tcgaatttga ttctgtgcga 8460
tagcgcccct gtgtgttctc gttatgttga ggaaaaaaat aatggttgct aagagattcg 8520
aactcttgca tcttacgata cctgagtatt cccacagtta actgcggtca agatatttct 8580
tgaatcaggc gccttagacc gctcgccaa acaaccaatt acttggtgag aaatagagta 8640
taattatcct ataaatataa cgtttttgaa cacacatgaa caaggaagta caggacaatt 8700
gattttgaag agaattgtgga ttttgatgta attggtggga ttccattttt aataaggcaa 8760
taatattagg tatgtagata tactagaagt tctcctcgac 8800

<210> 30

<211> 10042

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:pREPxr-CRHR

<400> 30

aagcttgcac gcctgcaggc cgatcgactc tagaggatca gaaaattatc gccataaaaag 60
acagaataag tcatcagcgg ttgtttcatt tcctatatct tttttttatt tttttatttt 120
ttaataaggg aaaatttaac gtctaaggat acagaagatt gtagcacat taaagtaata 180
aaggcttaag tagtaagtgc cttagcatgt tattgtattt caaaggacat aatctaaaat 240
aataacaata tcatttctca caagttattc aattttcttt tttttttcta ataatatcaa 300
gaatgtatta ttgttttgac ataagtcaac taatttattt aatatgctgg attaattctg 360
cagacatgta aattaacaag ttttagtcaa ataacgttga agtttcaatg aactcaaata 420
atttctcttt ttttttatat aaccatatgt ctaatctgat ttatatcttc cgcagggatc 480
aactgaagtt atgacatttg gattggatca cttataacct tggtcgcaa ataatacaaa 540
aatcagcgtt ataaaacaaa gaaggttttt gttaagaaat taatcctctt tcttgataag 600
aaagttgaac cgaaattgca gatactgata tatgaaaata ataccacaa ttttggaat 660
agcgcaagcc tcaatttaaa caatagggtga ggacacatga taatgacctc aatgattgtt 720
agaagaaaag agcctcatta caaaatcgaa aaatgaatgg ttgggtacaa gtttccaaaa 780
catggtaaag tggactttgc gtatgagacg taaatagaaa aaaacacttg ttatatgttt 840
tctagaatta ttgttgtctc tttatggttg gatgatgcaa aatagtaatt tcggttagtt 900
gctgtaaaac accacgagac aaatagatat ggatatttat taaatcagga aaaacgtaac 960
tctcggctac tggatggttc agtcaccaa cgattactgg ggagagaaaa cagggcaaaa 1020
gcaaagctta aaggaatccg attgtcattc ggcaatgtgc agcgaaacta aaaaccggat 1080
aatggacctg ttaatcgaaa cattgaagat atataaagga agaggaatcc tggcatatca 1140
tcaattgaat aagttgaatt aattatttca atctcattct cactttctga cttatagtcg 1200
ctttgttaaa ttggcctcga ggtcgactct agaattggag ggcaccgcga gctccgtctc 1260
gtcaaggccc ttctccttct ggggctgaac cccgtctctg cctccctcca ggaccagcac 1320

tgcgagagcc tgtccctggc cagcaacatc tcaggactgc agtgcaacgc atccgtggac 1380
ctcattggca cctgctggcc ccgcagccct gcggggcagc tagtggttcg gccctgccct 1440
gcctttttct atggtgtccg ctacaatacc acaaacaatg gctaccggga gtgcctggcc 1500
aatggcagct gggccgccc cgtgaattac tccgagtgc aggagatcct caatgaggag 1560
aaaaaaagca aggtgacta ccatgtcgca gtcacatca actacctggg ccaactgtatc 1620
tccctgggtg ccctcctggg ggcctttgtc ctctttctgc ggctcaggag catccgggtg 1680
ctgcgaaaca tcatccactg gaacctcact tccgccttca tccctgcgca cggccacctg 1740
ttcgtgggtc agctaaccat gagccccgag gtccaccaga gcaacgtggg ctggtgcagg 1800
ttggtgacag ccgcctacaa ctacttccat gtgaccaact tcttctggat gttcggcgag 1860
ggctgctacc tgcacacagc catcgtgctc acctactcca ctgaccggct gcgcaaattg 1920
atgttcatct gcattggctg ggggtgtgcc ttcccatca ttgtggcctg ggccattggg 1980
aagctgtact acgacaatga gaagtgtgg tttggcaaaa ggcctggggg gtacaccgac 2040
tacatctacc agggccccc gatcctgggtc ctgctgatca atttcatctt cttttcaac 2100
atcgtccgca tctcatgac caagctccgg gcatccacca cgtctgagac cattcagtac 2160
aggaaggctg tgaaagccac tctggtgctg ctgcccctcc tgggcatcac ctacatgctg 2220
ttcttcgtca atcccgggga ggatgaggtc tcccgggtcg tcttcatcta cttcaactcc 2280
ttcctggaat ccttccaggg cttctttgtg tctgtgttct actgtttcct caatagttag 2340
gtccgttctg ccatccggaa gaggtggcac cgggtggcagg acaagcactc gatccgtgcc 2400
cgagtggccc gtgccatgtc catccccacc tcccacccc gtgtcagctt tcacagcatc 2460
aagcagtcca cagcagtctg aggatccccg ggtaaaagga atgtctccct tgccagtact 2520
gctagggttt ttctttcaaa ctatggaagc ccattcaagc tgcataattac gattttgttt 2580
ttcgttttta gaaagtgggt tagatgagat aatagaaaaa ttcttgatct ccgacaacga 2640
gtacttttat ttttttgtc aatcacttta ctcaatatta gctcgaatc gtagaaacgt 2700
agacgggtgc gggataccga gtggtgtagt taagaatttt tataaaccac gtggcccaaa 2760
aatatgaacc caaacgttt atacatgagt atactttaag aaggctatac cccttcgtgt 2820
tagatgtagt ttagctacc caaccgagt ctatgagctt gacttcagat gtagaaggca 2880
ttaaatcgtt ttgaatatta attaaaaaac gatgaaaatt aaatatttaa aagcaatcat 2940
acgtgaaaa tttagtgtc tggctaattc ttcaacatgg aaatgccata aaagtgactt 3000
tgacaaaaaa aaaagtatat acaggtagta aactcatcta cttcattgac tttgtttaca 3060
gcatgtggaa ggaggaatat ttattgctaa atcgtagttt aacattcaat aagtaatact 3120
attgaaattc gacaagattg gccgcatgga tgaaaaagag gcattttgtt ttgggagaat 3180
tagttcaaat tagaactgaa aaaaaaaact ttacgaggca aaaatgtcgg attgagatcg 3240
taaaagtctg ctgctgctt tttgctttgt gattgttttc atggatacat cttgctggat 3300
atttaaaatt tagtactatg tataagatat tctataaatg ttttatcacc caaacctgtt 3360
agcgccttct taattctatt caatctggct tttgctctga gactacttct tggactttca 3420
ctacttggtt gttatacggg atttgtgtta ttagaagtga aataatcctt tctattagta 3480
atgcgagctc gaattcgagt ctaactcctt aaccactcgg acatagtac ttatctgaca 3540
ctcattgtaa attaatatat atataggcat tttgtttagt taaaggtagt taagtaatta 3600
gtataaacga accaatttta taatcaggaa gttaagtga ttgtagcaca tgtcgtaaaa 3660
attgtgaatt tttattgaat aatattttta atacaagcct ttctagacta ggtataactca 3720
taaacatata tgagcaaaaag gatagaggag attacattgc atcttctaca aatttattta 3780
ttgcccttta ctgaaaaatt aaataatgag tactaaatga taaaaagcgc tcagtacaag 3840
aaagattgca aaaatattgc attcttcagt aattaaagtt gcatataagg catattgaaa 3900
gtaatagtac taaaacagca gttagcgaat attaatagaa ttatattcgc aagacaattg 3960
tgacattaaa ttaaaaaatt gtagaatttt tactatcctc ttaacgcca tgagccttta 4020
taaaaagggt aaattagttt taacattctt tttttgagta agagtttaca atttatcaaa 4080
acctgtgtta ttatattcat tagtttcaat ttattagcat ctagagaaaa atcaattggc 4140
agttacattg ttggaattta tgaagaaaaa gaatctacaa cggagaataa gttgctgatc 4200

gctttcccta aaattgtata ttttgctgag cttattttga catttcgttg aagttttctc 4260
taattcgcat tcattttaag taaaacaatg agaaataaaa ttacaaaaaa tacaaattaa 4320
aatacaatth ttagctataa ttatagacga tgcctttgta tccattcttg tctcgcttgc 4380
ccctactttt tatcttttat ataccataat gaacgctgcc gctactaacc ataccccgat 4440
tttacatttc ggactcccaa ggacgtacaa aatagaaaac tatagaaaaa aataatcaga 4500
aaatagcatg tcatctcttt gtaaaacgag tttgcaagaa gaaaggaaac aatggagaag 4560
agatcatcca tttgtatgta aatttttagta aacttgaaga aatcactaac aacttctctt 4620
acttagggat tctatgcaaa accttgtaaa tcatctgatg gaggactcga tttaatgaat 4680
tggaaggttg gaattcactg gccgtcgttt tacaacgtcg tgactgggaa aaccctggcg 4740
ttaccaact taatcgccct gcagcacatc cccctttcgc cagctggcgt aatagcgaag 4800
aggcccgcac cgatcgccct tcccaacagt tgcgcagcct gaatggcgaa tggcgccgta 4860
tgcggtatth tctccttacg catctgtgag gtatttcaca ccgcatacgt caaagcaacc 4920
atagtacgag cctgttagcg gcgcattaag cgcggcggtt gtggtggtta cgcgcagcgt 4980
gacgcctaca ctggccagcg ccttagcgcc cgtcccttcc gctttcttcc ctctcttctt 5040
cgccacgttc gccggcttcc cccgtcaagc tctaaatcgg gggctccctt tagggttccg 5100
atthagtgct ttacggcacc tcgaccccaa aaaacttgat ttgggtgatg gttcacgtag 5160
tgggacatcg cctgataga cgtttttcgc ccttttgacg ttggagtcca cgttctttaa 5220
tagtggactc ttgttccaaa ctggaacaac actcaactct atctcgggct attcttttga 5280
tttataaggg attttgccga tttcgggtcta ttggttaaaa aatgagctga tttacaaaaa 5340
atttaacgag aattttaaca aaatatatac gtttacaatt ttatggtgca ctctcagtac 5400
aatctgctct gatgccgat agttaagcca gcccgacac ccgccaacac ccgctgacgc 5460
gccctgacgg gcttgtctgc tcccgccatc cgtttacaga caagctgtga ccgtctccgg 5520
gagctgcatg tgtcagaggt tttcacgctc atcacgaaa cgcgcgagac gaaaggccct 5580
cgtgatacgc ctatttttat aggttaatgt catgataata atggtttctt agacgtcagg 5640
tggaactttt cggggaaatg tgcgcggaac cctattttgt ttatttttct aaatacatte 5700
aaatatgtat ccgctcatga gacaataacc ctgataaatg cttcaataat attgaaaaag 5760
gaagagtatg agtattcaac atttccgtgt cgcctttatt cctttttttg cggcattttg 5820
ccttctgtt tttgctcacc cagaaacgct ggtgaaagta aaagatgctg aagatcagtt 5880
gggtgcacga gtgggttaca tcgaactgga tctcaacagc ggtaagatcc ttgagagttt 5940
tcgccccgaa gaacgttttc caatgatgag cactttttaa gttctgctat gtggcgcggt 6000
attatccgct attgacgccc ggcaagagca actcggctgc cgcatacact attctcagaa 6060
tgacttggtt gactactcac cagtacaga aaagcatctt acggatggca tgacagtaag 6120
agaattatgc agtctgcca taaccatgag tgataacact gcggccaact tacttctgac 6180
aacgatcgga ggaccgaagg agctaaccgc ttttttgac aacatggggg atcatgtaac 6240
tcgccttgat cgttggggaa cggagctgaa tgaagccata ccaaacgacg agcgtgacac 6300
cacgatgcct gtagcaatgg caacaacgtt gcgcaaaacta ttaactggcg aactacttac 6360
tctagcttcc cggcaacaat taatagactg gatggaggcg gataaagttg caggaccact 6420
tctgcgctcg gcccttccgg ctggctggtt tattgctgat aaatctggag ccggtgagcg 6480
tggtctcgc ggtatcattg cagcactggg gccagatggt aagccctccc gtatcgtagt 6540
tatctacagc acggggagtc aggcaactat ggatgaacga aatagacaga tcgctgagat 6600
aggtgcctca ctgattaagc attggttaact gtcagaccaa gtttactcat atatacttta 6660
gattgattta aaacttcatt ttttaatttaa aaggatctag gtgaagatcc tttttgataa 6720
tctcatgacc aaaatccctt aacgtgagtt ttcgtttcac tgagcgtcag accccgtaga 6780
aaagatcaaa ggatcttctt gagatccttt ttttctgcgc gtaatctgct gcttgcaaac 6840
aaaaaaacca ccgctaccag cgggtggtttg tttgccggat caagagctac caactctttt 6900
tccgaaggta actggcttca gcagagcgca gataccaaat actgtccttc tagtgtagcc 6960
gtagttaggc caccattca agaactctgt agcaccgcct acatacctcg ctctgctaata 7020
cctgttacca gtggctgctg ccagtgccga taagtctgtt cttaccgggt tggactcaag 7080

acgatatgta ccgataagg cgcagcggtc gggctgaacg gggggttcgt gcacacagcc 7140
cagcttgag cgaacgacct acaccgaact gagataccta cagcgtgagc tatgagaaaag 7200
cgccacgctt cccgaaggga gaaaggcgga caggtatccg gtaagcggca gggtcggaac 7260
aggagagcgc acgagggagc ttccaggggg aaacgcctgg tatctttata gtcctgtcgg 7320
gtttcgccac ctctgacttg agcgtcgatt tttgtgatgc tcgtcagggg ggcggagcct 7380
atggaaaaac gccagcaacg cggccttttt acggttcctg gccttttgct ggccttttg 7440
tcacatgttc tttcctgctg tatccctga ttctgtggat aaccgtatta cgcctttga 7500
gtgagctgat accgctcgcc gcagccgaac gaccgagcgc agcagagtcag tgagcgagga 7560
agcggaagag cgccaatac gcaaaccgcc tctccccgcg cgttggccga ttcattaatg 7620
cagctggcac gacaggtttc ccgactggaa agcgggcagt gagcgcaacg caattaatgt 7680
gagttagctc actcattagg caccacaggc ttacacttt atgcttccgg ctcgatgtt 7740
gtgtggaatt gtgagcgat aacaatttca cacaggaaac agctatgacc atgattacgc 7800
caagcttgct gatcgactac gtcgttaagg ccgtttctga cagagtaaaa ttcttgaggg 7860
aactttcacc attatgggaa atggttcaag aaggtattga cttaaactcc atcaaagggt 7920
caggtcattg agtggttttt atttggtgta ttttttttt tttagagaaa atcctccaat 7980
atcaaattag gaatcgtagt ttcattgatt tctgttacac ctaacttttt gtgtgggtgcc 8040
ctcctccttg tcaatattaa tgttaaagt caattotttt tccttatcac gttgagccat 8100
tagtatcaat ttgcttacct gtattccttt actatcctcc tttttctcct tcttgataaa 8160
tgtatgtaga ttgcgtatat agtttcgtct accctatgaa catattccat tttgtaattt 8220
cgtgtcgttt ctattatgaa tttcatttat aaagtttatg taaaaaatc ataaaaaag 8280
agaatctttt taagcaagga ttttcttaac ttcttcggcg acagcatcac cgacttcggg 8340
ggtactgttg gaaccaccta aatcaccagt tctgatacct gcatccaaaa cctttttaac 8400
tgcactttca atggccttac cttcttcagg caagttcaat gacaatttca acatcattgc 8460
agcagacaag atagtggcga tagggtcaac cttattcttt ggcaaactct gagcagaacc 8520
gtggcatggt tcgtacaaac caaatgcggt gttcttgtct ggcaaagagg ccaaggacgc 8580
agatggcaac aaaccaagg aacctgggat aacggaggct tcatcggaga tgatgtcacc 8640
aaacatggtg ctggtgatta taataccatt taggtgggtt ggggttctta ctaggatcat 8700
ggcggcagaa tcaatcaatt gatgttgaac cttcaatgta gggaattcgt tcttgatggt 8760
ttcctccaca gtttttctcc ataactttga agaggccaaa acattagctt tatccaagga 8820
ccaaataggc aatggtggct catgttgtag ggccatgaaa gcggccattc ttgtgattct 8880
ttgcacttct ggaacggtgt attgttcaat atcccaagcg acaccatcac catcgtcttc 8940
ctttctctta ccaaagtaaa tacctcccac taattctctg acaacaacga agtcagtacc 9000
tttagcaaat tgtggcttga ttggagataa gtctaaaaga gagtcggatg caaagttaca 9060
tggtcttaag ttggcgtaca attgaagttc ttacggatt tttagtaaac cttgttcagg 9120
tctaactacta ccggtacccc atttaggacc acccacagca cctaacaaaa cggcatcaac 9180
cttcttgag gcttcagcg cctcatctgg aagtgggaca cctgtagcat cgatagcagc 9240
accaccaatt aaatgatttt cgaaatcgaa cttgacattg gaacgaacat cagaaatagc 9300
tttaagaacc ttaatggctt cggctgtgat ttcttgacca acgtgggtcac ctggcaaaac 9360
gacgatcttc ttaggggcag acataggggc agacattaga atggtatatc cttgaaatat 9420
atatatatat tgctgaaatg taaaaggtaa gaaaagttag aaagtaagac gattgctaac 9480
cacctattgg aaaaaacaat aggtccttaa ataatttgt caacttcaag tattgtgatg 9540
caagcattta gtcatgaacg cttctctatt ctatatgaaa agccggttcc ggcctctcac 9600
ctttcctttt tctcccaatt tttcagttga aaaaggtata tgctcagggc gacctctgaa 9660
attaacaaaa aatttccagt catcgaattt gattctgtgc gatagcgccc ctgtgtgttc 9720
tcgttatgtt gaggaaaaaa ataatggttg ctaagagatt cgaactcttg catcttacga 9780
tacctgagta ttcccacagt taactgcggt caagatattt cttgaatcag gcgccttaga 9840
ccgctcggcc aaacaaccaa ttactgttg agaaatagag tataattatc ctataaatat 9900
aacgtttttg aacacacatg aacaaggaag tacaggacaa ttgattttga agagaatgtg 9960

gattttgatg taattgttgg gattccattt ttaataaggc aataatatta ggtatgtaga 10020
tatactagaa gttctcctcg ac 10042

<210> 31

<211> 5446

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:pSP72 sxa2

<400> 31

gaactcgagc agctgaagct tgcattgcctg caggctcgact ctagaggatc ctataatggg 60
cacttctctt gatccttacc gtcataattgg ctgtactggg agtccttctg gttcacataa 120
tttaatttgg atgactgtct acaaagacaa actccgccgt tgcctgaat gcggatctgt 180
atacaaatga aaattcatgg gagatcccaa cgctgaacac agtcattagt atgcttacca 240
atcttccata tccccttgct attttcatac tcttttataa caatttagct cttttatttg 300
aactttgaaa aatcaatgaa aactgtatgc tgatctatct cttatttttg gaacgattcg 360
attattcgta aacgtatatc ttataaaata ttctatctat tcaaccatat cttagttgtt 420
aattcaagca ccattctttt tttctttag tagatcgcaa atatgatcta attatcggtc 480
cgagtttcat aaacgatagt tatttattga tataccacat cagtggtaag tcaataaaact 540
tgtattttta attttccgat attaaacttt taataaagca acgtattttc aagccattct 600
ctaaccattt gaggggtctt agatggagcc atgtgcccaa ccgagttgga caatgtaaag 660
gctaagttac gttcatccaa agtaaatcca ttagttgtct ctaaaacttc aggagattga 720
gtgaatccct gccatccatt ccatgttgta ttttgcaatg ccaaaagagt tccgggtccaa 780
agaatttgaa gatccaatgc cccagctaag aaagacactt tgtatttttc tgtaagcgt 840
ggaattattt caacgagaac tgattctacg ttgttactca caatctttt gtataagtca 900
aaattgcatc cgtctgcgaa gacaccctcg ccgcttgta acgcagttga agctttcgta 960
gcatgtaaag aagagcgaac atcttctctg ttcaggtatg tgattaaggg atcattatac 1020
tcattgaaac tacaatcaag gctaacaatca taagttatga cacagctgtt ctctagataa 1080
agtaaaaagt tagaaaggga atacaagtca catccagaaa ttgaattaaa tacattccca 1140
atgtcttcac catcgagagc ttcccgcttc cttagagaag tagatgtgct aaagttgtac 1200
tctgggtctc atattgggta ttgttcagta ggaaacgtca atctgttcaa aactgaatca 1260
tattgacatt ctttattgag ctttttaaac tcttcagata ttgtggaaga agtattatta 1320
aaatagtatc caagtttgct gatattgtcc acccaaatag aagcagtaat ttgttcttga 1380
gtttcgtagt ccgccgtaag tccactgacg attcccacgc ccataaaatt tatattcaaa 1440
gatggttcag aaagtaatgc ctctgcaaaa ttggcgctcc atatgctacc gtagctttcg 1500
ccgacaaggt agagcttttt cttcattaag tggggaaaact tttggtaaaa tgatttcagg 1560
gcatttacia aatctgaaga agcttcttca atcgtagttg tatatgccgc ctgtccttgg 1620
gagtaccccg taccaaatgg ctgatctaac cataacatgt tggcgaagtt tgtccaagat 1680
tcaggattaa gggatgggtga ggggctactt tgggatattt caatgggtcc attttcactg 1740
aaaaagccta atgttccagc gcatccaggt cctccttgaa gccatactat gaaagtttcc 1800
gaatcaacia cagcagggggc atatgtataa aaaagtgatt tatcggaatt cgcttccaag 1860
tatccagaat ataattccgg taatgatccc tttaattcag gaagtgattt aattcgatcc 1920
gaagaaggag cagcgtgttg tctaggttga acagtctgat tcgaggaagc agaactcgtg 1980
ttggcttgtt gtatactgca tttccaatgt acagtgtgaa ttggtaaagc gtgaatgata 2040
gtcaattcaa ttattataat agcaaaaaga gatttcaaaa ataaagacag cattgaaaag 2100

agagacaatg attttgagcc tatgagaaaa gcgttagcgc taaattaaat ataatattta 2160
tcataatttg aatatattga tgtattcagt ttcgaaacaa ttcttgaaaa gtgactataa 2220
ttttatacgc ttcagtattt cgaaaaatgt gaaaatgcta tgacttactc gaataaagaa 2280
accggattgt ttatctgatt ctttttcata agagttgcat gctatatatg ataatacata 2340
ataatgggga ttcacttgaa tagattagca ttgtccattg tttacaatca acaacaatag 2400
agatgggcaa caaagacacc cagcgaagta ccatttacta gtaataggct aattcattta 2460
agtttttcaa cacaatagtc cttaatgaaa aaaccaattg tttcgtttac gcggcaagtg 2520
aatagggtag gactagaata acctatggtc tgtttaaatc gaaaaccagc ttatttcagc 2580
agcgatcatt tatgcattct taatgcttaa caatgcgagg cgactcaaag aatggagaaa 2640
attgattgtt ttatgaccgg tggaatcgaa taatttgact tttgctatga aggcttttagc 2700
ccttagaaat aaagaaataa aataaaaaat agatagataa caagaaaaaa tcaatgtcaa 2760
ataatttagt tgtattaatg aaaaataaga aataactaaa ttaaaaaaat ataaatacat 2820
ttatatatta gtaacactat ttctttttaa agaccttgaa atattatgtg agaaacttaa 2880
aaactaataa aaataaataa atataaaaaat ataaagaaaa agaaaaaaat taaaaattat 2940
atatacaata tactaataaa catagtgatt aattaaatga aaaaaataaa tgtttcttaa 3000
agcatggtta aaacacacaa tggcggattt ctagccatgg taccgagctc gaattcatcg 3060
atgatatcag atctgcgggt ctccctatag tgagtcgtat taatttcgat aagccagggt 3120
aacctgcatt aatgaatcgg ccaacgcgcg gggagaggcg gtttgcgat tgggcgctct 3180
tcgccttctt cgctcactga ctgcgtgcgc tcggtcggtc ggctgcggcg agcggtatca 3240
gctcactcaa aggcggtaat acggttatcc acagaatcag gggataacgc aggaaagaac 3300
atgtgagcaa aaggccagca aaaggccagg aaccgtaaaa aggccgcgtt gctggcggtt 3360
ttccataggc tcgcgcctcc tgacgagcat cacaaaaatc gacgctcaag tcagagggtg 3420
cgaaaccgga caggactata aagataccag gcgtttcccc ctggaagctc cctcgtgcgc 3480
tctcctgttc cgaccctgcc gcttaccgga tactgtccg cctttctccc ttcgggaagc 3540
gtggcgcttt ctcatagctc acgctgtagg tatctcagtt cgggtgtaggt cgctcgtcc 3600
aagctgggct gtgtgcacga acccccgtt cagcccgacc gctgcgcctt atccggtaac 3660
tatcgtcttg agtccaaccc ggtaagacac gacttatcgc cactggcagc agccactggt 3720
aacaggatta gcagagcgag gtatgtaggc ggtgctacag agttcttgaa gtggtggcct 3780
aactacggct acactagaag aacagtattt ggtatctgcg ctctgctgaa gccagtacc 3840
ttcggaaaaa gagttggtag ctcttgatcc ggcaaacaaa ccaccgctgg tagcgggtgt 3900
ttttttgttt gcaagcagca gattacgcgc agaaaaaaag gatctcaaga agatcctttg 3960
atcttttcta cggggtctga cgctcagtg aacgaaaact cacgttaagg gattttggtc 4020
atgagattat caaaaaggat cttcacctag atccttttaa attaaaaatg aagttttaa 4080
tcaatctaaa gtatatatga gtaaaacttg tctgacagtt accaatgctt aatcagtgag 4140
gcacctatct cagcgatctg tctatttcgt tcatccatag ttgcctgact ccccgctcg 4200
tagataacta cgatacggga gggcttacca tctggcccca gtgctgcaat gataccgcga 4260
gaccacgct caccggctcc agatttatca gcaataaacc agccagccgg aaggcccgag 4320
cgcagaagtg gtcctgcaac tttatccgcc tccatccagt ctattaattg ttgccgggaa 4380
gctagagtaa gtagttcgcc agttaatagt ttgcgcaacg ttgttgccat tgctacaggc 4440
atcgtggtgt cacgctcgtc gtttggtatg gcttcattca gctccggttc ccaacgatca 4500
aggcgagtta catgatcccc catggttggtc aaaaaagcgg ttagctcctt cggtcctccg 4560
atcgttgctc gaagtaagtt ggccgcagtg ttatcactca tggttatggc agcactgcat 4620
aattctctta ctgtcatgcc atccgtaaga tgcttttctg tgactggtga gtactcaacc 4680
aagtcattct gagaatagtg tatgcggcga ccgagttgct cttgcccggc gtcaatacgg 4740
gataataacc cgccacatag cagaacttta aaagtgtca tcattggaaa acgttcttcg 4800
gggcgaaaaac tctcaaggat cttaccgctg ttgagatcca gttcgatgta acccactcgt 4860
gcacccaact gatcttcagc atcttttact ttcaccagcg tttctgggtg agcaaaaaa 4920
ggaaggcaaa atgccgcaaa aaagggaata agggcgacac ggaaatgttg aatactcata 4980

ctcttccttt ttcaatatta ttgaagcatt tatcagggtt attgtctcat gagcggatac 5040
atatttgaat gtatttagaa aaataaacia ataggggttc cgcgcacatt tccccgaaaa 5100
gtgccacctg acgtctaaga aaccattatt atcatgacat taacctataa aaataggcgt 5160
atcacgagge cctttcgtct cgcgcgtttc ggtgatgacg gtgaaaacct ctgacacatg 5220
cagctcccgg agacggtcac agcttgtctg taagcggatg ccgggagcag acaagcccgt 5280
cagggcgctg cagcgggtgt tggcgggtgt cggggctggc ttaactatgc ggcacagag 5340
cagattgtac tgagagtga ccatatggac atattgtcgt tagaacgcgg ctacaattaa 5400
tacataacct tatgtatcat acacatacga tttagggtgac actata 5446

<210> 32

<211> 5704

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:pSP72

sxa2::ura4+

<400> 32

gaactcgagc agctgaagct tgcattgctg caggctcgact ctagaggatc ctataatggt 60
cacttctctt gatccttacc gtcattattg ctgtactggt agtccttctg gttcacataa 120
tttaatttgg atgactgtct acaaagacaa actccgccgt tgtcctgaat gcggatctgt 180
atacaaatga aaattcatgg gagatcccaa cgtgaacac agtcattagt atgcttacca 240
atcttccata tccccttgct attttcatac tcttttataa caatttagct cttttatttg 300
aactttgaaa aatcaatgaa aactgtatgc tgatctattt cttatttttg gaacgattcg 360
attattcgta aacgtatatc ttataaataa ttctatctat tcaaccatat cttagttgtt 420
aattcaagca ccattctttt tttctttagt tcatgcggaa atatgatcta attatcggtc 480
cgagtttcat aaacgatagt tatttattga tataccacat cagtggtaag tcaataaact 540
tgtattttta attttccgat attaaacttt taggatccac catgtagtga tattgacgaa 600
actttttgac atctaattta ttctgttcca acaccaatgt ttataaccaa gttttatctt 660
gtttgtctac atggtatttt acattcatct acatacatct ttcattggct ttgtacatag 720
ttatcattac aagtctaaaa aaattcactc ttttcttatt caatgtcaat ccaagagaaa 780
agattgtggt aatgtttagt gagcatgttt aataaattac tatagcaaat tactttttat 840
tccaagggtg tttatctata atagttaata ttttagtcgc tacataaaat tttaccaaag 900
agtacttgta tactaattct aaatgccttc tgacataaaa cgcctaggaa aacaaacgca 960
aacaaggcat cgactttttc aataaccaac caaaaaaatt ttacattagt ctttttttaa 1020
tgctgagaaa gtctttgctg atatgccttc caaccagctt ctctatatct cttggcttcg 1080
acaacaggat tacgaccagc tccatagact ccacgaccaa caatgatgat atcgctaccg 1140
cagtttataa tcaattcttc aggagtacga tattgctgtc ccagcccgctc tcttttaaca 1200
tccaagccga taccagggga catagttatg tagtcgcttt gaaggttagg aaatcgacga 1260
ccagctataa agccaaagca aaaatcggtg tgcttctcaa accattctaa ggttttctct 1320
gtgtaggaac cagtagccaa agagcctttg gaagacattt cagccaaaag caagagacca 1380
cgtcccaaag gtaaaccaac ttctttgagg ccttggtataa taccctcgcc tggcactgta 1440
tggcaatttg tgatatgagc ccaagaagca attttggtaca caccagatgc atattgtagc 1500
ttgacggtat ttccaatgtc tgcgaatttg cgatcctcaa agataagaaa acgatgcttt 1560
ttacctaaag ccaccagttt ttctaccata tcttggtcga aatcctcgac aacgtcaata 1620
tgtgtcttga taacacagac atagggtcca attttatcta ccaattctaa gatttcggat 1680

ttcttcgtca aatcgaccgc gactgacaag ttgctttgct tttcttccat caaagccaac 1740
aatcccttg caatgggatt tttcatcccc tcagctctag ctgaatagct ttgaaatact 1800
ctagcatcca taactttgct tttaaacctt taatttcgat ccaagcaaaa aagagggttct 1860
tggtaggaca atacggtaag aaaacacgac atgtgcagag atgccgacga agcatagtta 1920
aactgggatg gtaaaatcaa ttaagaatth ataaagacaa aattgtataa gtctctaaaa 1980
catcttaatt atacctcaca gaactatcta aaatatattc acaaagtgc aacattatca 2040
tgaaaaagaa ccattttaat ttaaagcaag ggcattaagg cttatttaca gaatttctta 2100
cttttgtaaa gattataagg ctgattatct ttttcacat gccaaaaatt acacaagata 2160
gaatggatgt ttgaaattaa acgtgagtat acaaacaat acactaggta aatcgaaaca 2220
tttttttctc cattaagtaa caaattccta tttagagaaa gaatgctgag tagattaaat 2280
aatctataca aactttttta acacaaatgc atacatatag ccagtgggat ttgtagctac 2340
atgggtggatc ctgaaaagag agacaatgat tttgagccta tgagaaaagc gttagcgcta 2400
aattaaatat aatatattatc atatttgga tatattgat tattcagttt cgaaacaatt 2460
cttgaaaagt gactataatt ttatacgctt cagtatttcg aaaaatgtga aaatgctatg 2520
acttactcga ataaagaaac cggattgttt atctgattca ttttcataag agttgcatgc 2580
tatatatgat aatacataat aatggggatt cacttgaata gattagcatt gtccattgtt 2640
tacaatcaac aacaatagag atgggcaaca aagacacca gcgaagtacc atttactagt 2700
aataggctaa ttcatttaag tttttcaaca caatagtcct taatgaaaaa accaattgtt 2760
tcgtttacgc ggcaagtga tagggtagga ctagaataac ctatggtctg tttaaatcga 2820
aaaccagctt atttcagcag cgatcattta tgcattctta atgcttaaca atgcgaggcg 2880
actcaaagaa tggagaaaat tgattgtttt atgaccgggtg gaatcgaata atttgacttt 2940
tgctatgaag gcttttagccc ttagaaataa agaaataaaa taaaaaatag atagataaca 3000
agaaaaaatc aatgtcaaat aatttagttg tattaatgaa aaataagaaa taactaaatt 3060
aaaaaaatat aaatacattt atatatagat aacactattt ctttttaag acctgaaat 3120
attatgtgag aaacttaaaa actaataaaa ataaataaat ataaaaatat aaagaaaaag 3180
aaaaaaatta aaaattatat atacaatata ctaataaaca tagtgattaa ttaaatgaaa 3240
aaaataaatg tttcttaaaag catggttaaa acacacaatg gcggatttct agccatggta 3300
ccgagctcga attcatcgat gatatcagat ctgccggtct ccctatagt agtcgtatta 3360
atttcgataa gccagggtta cctgcattaa tgaatcgcc aacgcgcggg gagaggcgg 3420
ttgcgtattg ggcgctcttc cgcttcctcg ctcaactgact cgctgcgctc ggtcgttcg 3480
ctgcggcgag cgggtatcag tcaactcaaag gcggtataac ggttatccac agaatacagg 3540
gataacgcag gaaagaacat gtgagcaaaa ggccagcaaa aggccaggaa ccgtaaaaag 3600
gccgcgttgc tggcggtttt ccataggctc cgccccctg acgagcatca caaaaatcga 3660
cgctcaagtc agaggtggcg aaaccgcaga ggactataaa gataccaggc gtttccccct 3720
ggaagctccc tcgtgcgctc tcctgttccg accctgccgc ttaccggata cctgtccgcc 3780
tttctccctt cgggaagcgt ggcgctttct catagctcac gctgtaggta tctcagttcg 3840
gtgtaggtcg ttcgctccaa gctgggctgt gtgcacgaac cccccgttca gccgaccgc 3900
tgcgcttat ccggttaacta tcgtcttgag tccaaccgg taagacacga cttatcgcca 3960
ctggcagcag cactggtaaa caggattagc agagcgagg atgtaggcg tgctacagag 4020
ttcttgaagt ggtggcctaa ctacggctac actagaagaa cagtatttgg tatctgcgct 4080
ctgctgaagc cagttacctt cggaataaga gttggtagct cttgatccgg caaacaacc 4140
accgctggta gcggtggttt tttgtttgc aagcagcaga ttacgcgcag aaaaaagga 4200
tctcaagaag atcctttgat cttttctac gggctctgac ctcaagtggaa cgaaaactca 4260
cgtaaggga ttttggtcat gagattatca aaaaggatct tcacctagat ctttttaaat 4320
taaaaatgaa gttttaaatc aatctaaagt atatatgagt aaacttggtc tgacagttac 4380
caatgcttaa tcagtgggc acctatctca gcgatctgtc tatttcgttc atccatagtt 4440
gcctgactcc ccgtcgtgta gataactacg atacgggag gcttaccatc tggccccagt 4500
gctgcaatga taccgcgaga ccacgctca ccggtccag atttatcagc aataaaccag 4560

ccagccggaa gggccgagcg cagaagtggc cctgcaactt tatccgcctc catccagtct 4620
attaattggt gccgggaagc tagagtaagt agttcgccag ttaatagttt gcgcaacgtt 4680
gttgccattg ctacaggcat cgtggtgtca cgctcgctgt ttggtatggc ttcattcagc 4740
tccggttccc aacgatcaag gcgagttaca tgatcccca tgttggtgcaa aaaagcgggt 4800
agctccttcg gtccctccgat cgttggtcaga agtaagttgg ccgcagtgtt atcactcatg 4860
gttatggcag cactgcataa ttctcttact gtcattgccat ccgtaagatg cttttctgtg 4920
actggtgagt actcaaccaa gtcattctga gaatagtgtg tgccggcgacc gagttgctct 4980
tgcccggtgt caatacggga taataccgcg ccacatagca gaactttaaa agtgctcatc 5040
attggaaaac gttcttcggg gcgaaaactc tcaaggatct taccgctgtt gagatccagt 5100
tcgatgtaac cactcgtgc acccaactga tcttcagcat cttttacttt caccagcgtt 5160
tctgggtgag caaaaacagg aaggcaaat gccgcaaaa agggaataag ggcgacacgg 5220
aaatgttgaa tactcatact ctctctttt caatattatt gaagcattta tcagggttat 5280
tgtctcatga gcggatacat atttgaatgt atttagaaaa ataaacaaat aggggttccg 5340
cgcacatttc cccgaaaagt gccacctgac gtctaagaaa ccattattat catgacatta 5400
acctataaaa ataggcgtat caccaggccc ttctgtctcg cgcgtttcgg tgatgacggg 5460
gaaaacctct gacacatgca gctcccgag acggtcacag cttgtctgta agcggatgcc 5520
gggagcagac aagcccgta gggcgcgta gcgggtgttg gcgggtgtcg gggctggctt 5580
aactatgcgg catcagagca gattgtactg agagtgcacc atatggacat attgtcgtta 5640
gaacgcggct acaattaata cataacctta tgtatcatat acatacgatt taggtgacac 5700
tata 5704

<210> 33

<211> 6952

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:pSP72 sxa>lacZ

<400> 33

tatagtgtca cctaaatcgt atgtgtatga tacataaggt tatgtattaa ttgtagccgc 60
gttctaacga caatatgtcc atatggtgca ctctcagtac aatctgctct gatgccgcat 120
agttaagcca gcccgcacac ccgccaacac ccgtgacgc gccctgacgg gcttgtctgc 180
tcccggcatc cgottacaga caagctgtga ccgtctccgg gagctgcatg tgtcagaggt 240
tttcaccgtc atcacgaaa cgcgcgagac gaaagggcct cgtgatacgc ctatttttat 300
aggtaaatgt catgataata atggtttctt agacgtcagg tggcactttt cggggaaatg 360
tgcgcggaac ccctatttgt ttatttttct aaatacatc aaatatgtat ccgctcatga 420
gacaataacc ctgataaatg cttcaataat attgaaaaag gaagagtatg agtattcaac 480
atttcctgtg cgcccttatt cccttttttg cggcattttg ccttctgtt tttgtctacc 540
cagaaacgct ggtgaaagta aaagatgctg aagatcagtt ggggtgcacga gtgggttaca 600
tcgaactgga tctcaacagc ggtaagatcc ttgagagttt tcgccccgaa gaacgttttc 660
caatgatgag cactttttaa gttctgctat gtggcgcggt attatcccg attgacgccg 720
ggcaagagca actcggtcgc cgcatacact attctcagaa tgacttggtt gactactcac 780
cagtcacaga aaagcatctt acggatggca tgacagtaag agaattatgc agtgctgcc 840
taaccatgag tgataaacact gcggccaact tacttctgac aacgatcgga ggaccgaagg 900
agctaaccgc ttttttgac aacatggggg atcatgtaac tcgccttgat cgttgggaac 960
cggagctgaa tgaagccata ccaaacgacg agcgtgacac cacgatgcct gtagcaatgg 1020

caacaacggt ggcgaaacta ttaactggcg aactacttac tctagcttcc cggcaacaat 1080
taatagactg gatggaggcg gataaagttg caggaccact tctgogctcg gcccttccgg 1140
ctggctggtt tattgctgat aaatctggag ccggtgagcg tgggtctcgc ggtatcattg 1200
cagcactggg gccagatggt aagccctccc gtatcgtagt tatctacacg acggggagtc 1260
aggcaactat ggatgaacga aatagacaga tcgctgagat aggtgcctca ctgattaagc 1320
attggttaact gtcagaccaa gtttactcat atatacttta gattgattta aaacttcatt 1380
tttaatttaa aaggatctag gtgaagatcc tttttgataa tctcatgacc aaaatccctt 1440
aacgtgagtt ttcgttccac tgagcgtcag accccgtaga aaagatcaaa ggatcttctt 1500
gagatccttt ttttctgccc gtaatctgct gcttgcaaac aaaaaaacca ccgctaccag 1560
cgggtggtttg tttgcccgat caagagctac caactctttt tccgaaggta actggcttca 1620
gcagagcgca gataccaaat actgttcttc tagtgtagcc gtagttaggc caccacttca 1680
agaactctgt agcaccgcct acatacctcg ctctgctaact cctgttacca gtggctgctg 1740
ccagtggcga taagtctgtt cttaccgggt tggactcaag acgatagtta ccggataagg 1800
cgcagcggtc gggctgaacg ggggggtcgt gcacacagcc cagcttggag cgaacgacct 1860
acaccgaact gagataccta cagcgtgagc tatgagaaag cgccacgctt cccgaaggga 1920
gaaaggcgga caggatccg gtaagcggca gggctcggaac aggagagcgc acgagggagc 1980
ttccaggggg aaacgcctgg tatctttata gtctgtcgcg gtttcgccac ctctgacttg 2040
agcgtcgatt tttgtgatgc tcgtcagggg ggcggagcct atggaaaaac gccagcaacg 2100
cggccttttt acggttccctg gccttttgct ggctttttgc tcacatgttc tttcctgcgt 2160
tatcccctga ttctgtggat aaccgtatta ccgcctttga gtgagctgat accgctcgcc 2220
gcagccgaac gaccgagcgc agcaggtcag tgagcgagga agcgggaagag cgccaatac 2280
gcaaaccgcc tctccccgcg cgttggccga ttcattaatg caggttaacc tggcttatcg 2340
aaattaatac gactcactat agggagaccg gcagatctga tatcatcgat gaattcgagc 2400
tcggtagcat ggctagaaat ccgccattgt gtgttttaac catgctttaa gaaacattta 2460
tttttttcat ttaattaatc actatgttta ttagtatatt gtatatataa tttttaattt 2520
ttttcttttt ctttatattt ttatatattt ttatttttat tagtttttaa gtttctcaca 2580
taatatttca aggtctttta aaagaaatag tggtactaat atataaatgt atttatattt 2640
ttttaattta gttatttctt atttttcatt aatacaacta aattatttga cattgatttt 2700
ttcttgttat ctatctattt tttattttat ttctttattt ctaagggcta aagccttcat 2760
agcaaaagtc aaattattcg attccaccgg tcataaaaca atcaattttc tccattcttt 2820
gagtcgcctc gcattgttaa gcattaagaa tgcataaatg atcgctgctg aaataagctg 2880
gttttcgatt taaacagacc ataggttatt ctagtcttac cctattcact tgccgcgtaa 2940
acgaaacaat tggttttttc attaaggact attgtgttga aaaacttaaa tgaattagcc 3000
tattactagt aaatggtact tcgctgggtg tctttgttgc ccatctctat tgttggtgat 3060
tgtaacaat ggacaatgct aatctattca agtgaatccc cattattatg tattatcata 3120
tatagcatgc aactcttatg aaaatgaatc agataaaca tccggtttct ttattcgagt 3180
aagtcatagc attttcacat ttttcgaaat actgaagcgt ataaaattat agtcactttt 3240
caagaattgt ttcgaaactg aatacatcaa tatattccaa atatgataaa tattatattt 3300
aatttagcgc taacgctttt ctcataggct caaaatcatt gtctctcttt tcaatgcagc 3360
tggcacgaca ggtttcccga cttaatcgcc ttgcagcaca tccccctttc gccagctggc 3420
gtaatagcga agaggcccgcc accgatcgcc cttcccaaca gttgcgcagc ctgaatggcg 3480
aatggcgctt tgcctgggtt ccggcaccag aagcgggtgcc ggaaagctgg ctggagtgcg 3540
atcttctctga ggccgatact gtctgctgcc cctcaaactg gcagatgcac ggttacgatg 3600
cgcccatcta caccaacgta acctatcca ttacgggtcaa tccgccgttt gttcccacgg 3660
agaatccgac gggttgttac tcgctcacat ttaatgttga tgaaagctgg ctacaggaag 3720
gccagacgcg aattattttt gatggcggtta actcggcggt tcatctgtgg tgcaacgggc 3780
gctgggtcgg ttacggccag gacagtcgtt tgccgtctga atttgacctg agcgcathtt 3840
tacgcgcggc agaaaaccgc ctgcgggtga tgggtgctgcg ttggagtgac ggcagttatc 3900

tggaagatca ggatatgtgg cggatgagcg gcattttccg tgacgtctcg ttgctgcata 3960
aaccgactac acaaatcagc gattttccatg ttgccactcg ctttaatgat gatttcagcc 4020
gcgctgtact ggaggtctgaa gttcagatgt gcggcgagtt gcgtgactac ctacgggtaa 4080
cagtttcttt atggcagggt gaaacgcagg tcgccagcgg caccgcgcct ttcggcgggtg 4140
aaattatcga tgagcgtggt ggttatgccg atcgcgtcac actacgtctg aacgtcgaaa 4200
accgaaaact gtggagcgcc gaaatcccga atctctatcg tcgggtggtt gaactgcaca 4260
ccgccgacgg cacgctgatt gaagcagaag cctgcgatgt cggtttccgc gaggtgcgga 4320
ttgaaaatgg tctgctgctg ctgaacggca agccgttgct gattcgaggc gttaaccgtc 4380
acgagcatca tcctctgcat ggtcaggta tggatgagca gacgatggtg caggatatcc 4440
tgctgatgaa gcagaacaac tttaacgccg tcgctgttcc gcattatccg aaccatccgc 4500
tgtggtacac gctgtgcgac cgctacggcc tgtatgtggt ggatgaagcc aatattgaaa 4560
cccacggcat ggtgccaatg aatcgtctga ccgatgatcc gcgctggcta ccggcgatga 4620
gcgaacgcgt aacgcgaatg gtgcagcgcg atcgtaatca cccgagtgtg atcatctggt 4680
cgctggggaa tgaatcaggc cacggcgcta atcacgacgc gctgtatcgc tggatcaaat 4740
ctgtcgatcc ttcccgcccg gtgcagtatg aaggcggcgg agccgacacc acggccaccg 4800
atattatttg cccgatgtac gcgcgcgtgg atgaagacca gcccttcccg gctgtgccga 4860
aatggtccat caaaaaatgg ctttcgctac ctggagagac gcgcccgtg atcctttgcg 4920
aatacgcca cgcatgggt aacagtcttg gcggtttcgc taaatactgg caggcgtttc 4980
gtcagtatcc ccgtttacag ggcggcttcg tctgggactg ggtggatcag tcgctgatta 5040
aatatgatga aaacggcaac ccgtggtcgg cttacggcgg tgattttggc gatacgcca 5100
acgatcgcca gttctgtatg aacggtctgg tctttgccga ccgcacgcc catccagcgc 5160
tgacggaagc aaaacaccag cagcagtttt tccagttccg tttatccggg caaaccatcg 5220
gaagtgaccag cgaataacctg ttccgtcata gcgataacga gctcctgcac tggatggtg 5280
cgctggatgg taagccgctg gcaagcgggt aagtgcctct ggatgtcgt ccacaaggta 5340
gaacagttgat tgaactgcct gaactaccgc agccggagag cgcgggcaa ctctggctca 5400
cagtacgcgt agtgcaaccg aacgcgaccg catggtcaga agccgggcac atcagcgct 5460
ggcagcagtg gcgtctggcg gaaaacctca gtgtgacgct ccccgcccg tccacgcca 5520
tcccgcatct gaccaccagc gaaatggatt tttgcatcga gctgggtaaat aagcgttggc 5580
aatttaaccg ccagtcaggc tttctttcac agatgtggat tggcgataaa aaacaaactgc 5640
tgacgccgct gcgcgatcag ttcaaccgtg caccgctgga taacgacatt ggcgtaagt 5700
aagcgaccgg cattgacct aacgcctggg tcgaacgctg gaaggcggcg ggcattacc 5760
aggccgaagc agcgttgttg cagtgcacgg cagatacact tgctgatcg gtgctgatta 5820
cgaccgctca cgcgtggcag catcagggga aaaccttatt tatcagccgg aaaacctacc 5880
ggattgatgg tagtggtaaa atggcgatta ccgttgatgt tgaagtggcg agcgatacac 5940
cgcatccggc gcggtattgg ctgaactgcc agctggcgca ggtagcagag cgggtaaaact 6000
ggctcggatt agggccgcaa gaaaactatc ccgaccgct tactgccgc tgttttgacc 6060
gctgggatct gccattgtca gacatgtata ccccgtagct ctccccgagc gaaaacggtc 6120
tgcgctgcgg gacgcgcgaa ttgaattatg gccacacca gtggcgcggc gacttccagt 6180
tcaacatcag ccgctacagt caacagcaac tgatggaaac cagccatcgc catctgctgc 6240
acgcggaaga aggcacatgg ctgaatatcg acggtttcca tatggggatt ggtggcgacg 6300
actcctggag ccgctcagta tcggcggaat tccagctgag cgcgggtcgc taccattacc 6360
agttggtctg gtgtcaaaaa taaaagttta atatcggaaa atttaaaata caagtttatt 6420
gacttaccac tgatgtggtat tatcaataaa taactatcgt ttatgaaact cggaaacgata 6480
attagatcat atttccgcat gactacaaga aaaaaagaat ggtgcttgaa ttaacaacta 6540
agatatggtt gaatagatag aattatttat aagatatacg ttacgaata atcgaatcgt 6600
tcccaaaaata agaaatagat cagcatacag ttttcattga tttttcaaag ttcaataaaa 6660
agagctaaat tggtataaaa gagtatgaaa atagcaaggg gatatggaag attggtgaagc 6720
ataactaatga ctgtgttcag cggtgggatc tcccatgaat ttttaatttg atacagatcc 6780

gcattcagga caacggcgga gtttgtcttt gtagacagtc atccaaatta aattatgtga 6840
accagaagga ctaccagtac agccaatatg acggtaagga tcaagagaag tgaccattat 6900
aggatcctct agagtcgacc tgcaggcatg caagcttcag ctgctcgagt tc 6952

<210> 34

<211> 4717

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:pSP72 sxa2>ura4

<400> 34

tatagtgtca cctaaatcgt atgtgtatga tacataaggt tatgtattaa ttgtagccgc 60
gttctaacga caatatgtcc atatggtgca ctctcagtac aatctgctct gatgccgcat 120
agttaagcca gccccgacac ccgccaacac ccgctgacgc gccctgacgg gcttgtctgc 180
tcccggcatc cgcttacaga caagctgtga ccgtctccgg gagctgcatg tgtcagaggt 240
tttcaccgtc atcaccgaaa cgcgcgagac gaaagggcct cgtgatacgc ctatTTTTAT 300
aggTTaatgt catgataata atggTTTTctt agacgtcagg tggcactttt cggggaaatg 360
tgcgcggaac ccctatttgt ttatTTTTctt aaatacattc aaatatgtat ccgctcatga 420
gacaataacc ctgataaatg cttcaataat attgaaaaag gaagagtatg agtattcaac 480
atttccgtgt cgcccttatt ccctTTTTtg cggcattttg ccttcctggt tttgctcacc 540
cagaaaacgct ggtgaaagta aaagatgctg aagatcagtt gggTgcacga gtgggttaca 600
tcgaactgga tctcaacagc ggtaagatcc ttgagagttt tcgccccgaa gaacgttttc 660
caatgatgag cactttttaa gttctgctat gtggcgcggt attatcccggt attgacgccg 720
ggcaagagca actcggtcgc cgcatacact attctcagaa tgacttggtt gagtactcac 780
cagtcacaga aaagcatctt acggatggca tgacagtaag agaattatgc agtgctgcca 840
taaccatgag tgataaacact gcggccaact tacttctgac aacgatcgga ggaccgaagg 900
agctaaccgc ttttttgcac aacatggggg atcatgtaac tcgccttgat cgttgggaaac 960
cggagctgaa tgaagccata ccaaacgacg agcgtgacac cacgatgcct gtagcaatgg 1020
caacaacggt gcgcaaaacta ttaactggcg aactacttac tctagcttcc cggcaacaat 1080
taatagactg gatggaggcg gataaagttg caggaccact tctgcgctcg gcccttcogg 1140
ctggctgggt tattgtgat aaatctggag ccggtgagcg tgggtctcgc ggtatcattg 1200
cagcactggg gccagatggt aagccctccc gtatcgtagt tatctacacg acggggagtc 1260
aggcaactat ggatgaacga aatagacaga tcgctgagat aggtgcctca ctgattaagc 1320
attggttaact gtcagaccaa gtttactcat atatacttta gattgattta aaacttcatt 1380
tttaatttaa aaggatctag gtgaagatcc tttttgataa tctcatgacc aaaatccctt 1440
aacgtgagtt ttcgttccac tgagcgtcag accccgtaga aaagatcaaa ggatcttctt 1500
gagatccctt ttttctgcgc gtaatctgct gcttgcaaac aaaaaaacca ccgctaccag 1560
cggTggtttg tttgccggat caagagctac caactctttt tccgaaggta actggcttca 1620
gcagagcgca gataccaaat actgttcttc tagtgtagcc gtagttaggc caccacttca 1680
agaactctgt agcaccgcct acatacctcg ctctgctaata cctgttacca gtggctgctg 1740
ccagtggcga taagtctgtg cttaccgggt tggactcaag acgatagtta ccggataagg 1800
cgcagcggtc gggctgaacg ggggggttcgt gcacacagcc cagcttgagg cgaacgacct 1860
acaccgaact gagataccta cagcgtgagc tatgagaaag cgccacgctt cccgaaggga 1920
gaaaggcgga caggtatccg gtaagcggca gggTcggaac aggagagcgc acgagggagc 1980
ttccaggggg aaacgcctgg tatctttata gtcctgtcgg gtttcgccac ctctgacttg 2040

```

agcgtcgatt tttgtgatgc tcgtcagggg ggcggagcct atggaaaaac gccagcaacg 2100
cggccttttt acggttcctg gccttttgct ggccttttgc tcacatgttc tttcctgcgt 2160
tatccctga ttctgtggat aaccgtatta ccgcctttga gtgagctgat accgctcgcc 2220
gcagccgaac gaccgagcgc agcagagtcag tgagcgagga agcggagag cgccaatac 2280
gcaaacccgcc tctccccgcg cgttggccga ttcattaatg caggttaacc tggcttatcg 2340
aaattaatac gactcactat agggagaccg gcagatctga tatcatcgat gaattcgagc 2400
tcggtagcat ggctagaaat ccgccattgt gtgttttaac catgctttaa gaaacattta 2460
tttttttcat ttaattaatc actatgttta ttagtatatt gtatatataa tttttaattt 2520
ttttcttttt ctttatattt ttatatattt ttatttttat tagtttttaa gtttctcaca 2580
taatatattca aggtctttta aaagaaatag tgttactaat atataaatgt atttatattt 2640
ttttaattta gttatttctt atttttcatt aatacaacta aattatttga cattgatttt 2700
ttcttggtat ctatctattt tttattttat ttctttattt ctaagggcta aagccttcac 2760
agcaaaagtc aaattattcg attccaccgg tcataaaaca atcaattttc tccattcttt 2820
gagtcgcctc gcattgttaa gcattaagaa tgcataaatg atcgctgctg aaataagctg 2880
gttttcgatt taaacagacc ataggttatt ctagtcctac cctattcact tgccgcgtaa 2940
acgaaacaat tggttttttc attaaggact attgtgttga aaaacttaaa tgaattagcc 3000
tattactagt aaatggtact tcgctgggtg tctttgttgc ccatctctat tgttggtgat 3060
tgtaaaacaat ggacaatgct aatctattca agtgaatccc cattattatg tattatcata 3120
tatagcatgc aactcttatg aaaatgaatc agataaaciaa tccggtttct ttattcgagt 3180
aagtcatagc attttcacat ttttcgaaat actgaagcgt ataaaattat agtcactttt 3240
caagaattgt ttcgaaactg aatacatcaa tatattccaa atatgataaa tattatattt 3300
aatttagcgc taacgctttt ctcataggct caaaatcatt gtctctcttt tcaatggatg 3360
ctagagtatt tcaaagctat tcagctagag ctgaggggat gaaaaatccc attgccaaag 3420
aattgttggc tttgatggaa gaaaagcaaa gcaacttgtc agtcgcggtc gatttgacga 3480
agaaatccga aatcttagaa ttggtagata aaattggacc ctatgtctgt gttatcaaga 3540
cacatattga cgttgtcgag gatttcgacc aggatatggt agaaaaactg gtggccttag 3600
gtaaaaagca tcgttttctt atctttgagg atcgcaaatt cgcagacatt ggaaataccg 3660
tcaagctaca atatgcatct ggtgtgtaca aaattgcttc ttgggctcat atcaciaaatt 3720
gccatacagt gccaggcgag ggtattatac aaggcctcaa agaagttggt ttacctttgg 3780
gacgtggtct cttgcttttg gctgaaatgt cttccaaagg ctctttggct actggttcct 3840
acacagagaa aaccttagaa tggtttgaga agcataccga tttttgcttt ggctttatag 3900
ctggtcgtcg atttcctaac cttcaaagcg actacataac tatgtcccct ggtatcggt 3960
tggtatgttaa aggagacggg ctgggacagc aatatcgtae tcctgaagaa gtgattgtaa 4020
actgcggtag cgatatcatc attgttggtc gtggagtcta tggagctggt cgtaatcctg 4080
ttgtcgaagc caagagatat agagaagctg gttggaaggc atatcagcaa agactttctc 4140
agcattaaaa gtttaatatc ggaaaattta aaatacaagt ttattgactt accactgatg 4200
tggtatatca ataaataact atcgtttatg aaactcgga cgataattag atcatatttc 4260
cgcatgacta caagaaaaaa agaatggtgc ttgaattaac aactaagata tggttgaata 4320
gatagaatta tttataagat atacgtttac gaataatcga atcgttcca aaataagaaa 4380
tagatcagca tacagttttt attgattttt caaagttcaa ataaaagagc taaattgtta 4440
taaaagagta tgaaaatagc aaggggatat ggaagattgg taagcatact aatgactgtg 4500
ttcagcggtg ggtctccca tgaattttta tttgtataca gatccgcat caggacaacg 4560
gcggagtttg tctttgtaga cagtcacca aattaaatta tgtgaaccag aaggactacc 4620
agtacagcca atatgacggt aaggatcaag agaagtgacc attataggat cctctagagt 4680
cgacctgcag gcatgcaagc ttcagctgct cgagttc 4717

```

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 35

ggggggtacc atggctagaa atccgccatt gtgtg

35

<210> 36

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 36

cttctcgtaa aggcacattg acgg

24

<210> 37

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 37

tgaaaagaga gacaatg

17

<210> 38

<211> 15

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer

used in the construction of the yeast and the G
alpha transplants

<400> 38
taaaagtтта atatc

15

<210> 39
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 39
atgcagctgg cacgacaggt ttcccgc

28

<210> 40
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 40
tttttgacac cagaccaact ggtaatggta gc

32

<210> 41
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 41
atggatgcta gagtatttc

19

<210> 42
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 42
atgctgagaa agtctttgc

19

<210> 43
<211> 32
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 43
tagattgttg gacataatcg tatcttgaac gg

32

<210> 44
<211> 35
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 44
gaatataatc ttgttttagat gaatttttcc ttaac

35

<210> 45
<211> 35
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 45

caatatgaac ttcttttagat gaatttttcc ttaac

35

<210> 46

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 46

ggatgcggac tttatttagat gaatttttcc ttaac

35

<210> 47

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 47

gattgcggac ttttttagat gaatttttcc ttaac

35

<210> 48

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 48

gaatgcggac tttattagat gaatttttcc ttaac

35

<210> 49

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 49

tatattggac ttgctagat gaatttttcc ttaac

35

<210> 50

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 50

gatattatgc ttcaatagat gaatttttcc ttaac

35

<210> 51

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer
used in the construction of the yeast and the G
alpha transplants

<400> 51

caacttatgc ttcaatagat gaatttttcc ttaac

35

<210> 52

<211> 35

<212> DNA

THIS PAGE BLANK (USPTO)